



05-21-01

PCT

JC19 Rec'd PCT/PTO 18 MAY 2001

FORM PCT/US (REV 10-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER P02196US0
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371			U.S. APPLICATION NO. (If known, see 37 CFR 1.5) <b>09/856339</b> <small>To be assigned</small>
INTERNATIONAL APPLICATION NO. PCT/GB99/03873	INTERNATIONAL FILING DATE 19 November 1999	PRIORITY DATE CLAIMED 19 November 1998	
TITLE OF INVENTION PROCESS FOR OXIDISING TERPENES			
APPLICANT(S) FOR DO/EO/US Luet Lok Wong; Stephen Graham Bell; and Angus Bishop Carmichael			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:			
<p>1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</p> <p>3. <input type="checkbox"/> This is an express request to promptly begin national examination procedures (35 U.S.C. 371 (f)).</p> <p>4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).</p> <p>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c)(2))</p> <p>a. <input checked="" type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau).</p> <p>b. <input type="checkbox"/> has been communicated by the International Bureau.</p> <p>c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371 (c)(2)).</p> <p>7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))</p> <p>a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau).</p> <p>b. <input type="checkbox"/> have been communicated by the International Bureau.</p> <p>c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</p> <p>d. <input type="checkbox"/> have not been made and will not be made.</p> <p>8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).</p> <p>9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).</p> <p>10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).</p>			
<b>Items 11 to 16 below concern document(s) or information included:</b>			
<p>11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</p> <p>12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 &amp; 3.31 is included.</p> <p>13. <input checked="" type="checkbox"/> A <b>FIRST</b> preliminary amendment.</p> <p><input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.</p> <p>14. <input type="checkbox"/> A substitute specification.</p> <p>15. <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>16. <input checked="" type="checkbox"/> Other items or information: (a) PCT Request; (b) PCT Demand; (c) PCT Receipt of Demand; and (d) PCT Notification of Change of Address for PCT Applicant</p>			

JG18 Rec'd PCT/PTO 1 8 MAY 2001

U.S. APPLICATION NO. (if known, see 37 CFR 1.5) <div style="font-size: 2em; font-weight: bold; margin-left: 50px;">09/856339</div> <div style="margin-left: 100px; font-size: 0.8em;">To be assigned</div>	INTERNATIONAL APPLICATION NO PCT/GB99/03873	ATTORNEY'S DOCKET NUMBER P02196US0
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------	---------------------------------------

17. ☒ The following fees are submitted:

**BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) – (5)):**

☐ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... \$1000.00

☒ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... \$860.00

☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... \$710.00

☐ International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... \$690.00

☐ International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) ..... \$100.00

**ENTER APPROPRIATE BASIC FEE AMOUNT =**

Surcharge of \$ 130.00 for furnishing the oath or declaration later than ☐ 20 ☒ 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	18 - 0 =	0	x 18.00	\$	0
Independent claims	3 - 0 =	0	x 80.00	\$	0
MULTIPLE DEPENDENT CLAIM(s) (if applicable)				x	\$ 0
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$	999.00
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	499.50
<b>SUBTOTAL =</b>				\$	499.50
Processing fee of \$ _____ for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)). +				\$	
<b>TOTAL NATIONAL FEE =</b>				\$	499.50
Fee for recording the enclosed assignment (37 CFR 1.21 (h)). Assignment must be accompanied by appropriate cover sheet (37 CFR 3.28, 3.31) ( _____ per property). +				\$	
<b>TOTAL FEES ENCLOSED =</b>				\$	499.50
				<b>Amount to be Refunded:</b>	\$
				<b>Charged:</b>	\$

a. ☐ A check in the amount of \$ \_\_\_\_\_ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 06-2375 in the amount of \$ 499.50 to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required or credit any overpayment to my Deposit Account No. 06-2375. A duplicate copy of this sheet is enclosed.

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO:

Thomas D. Paul  
 Fulbright & Jaworski LLP  
 1301 McKinney, Suite 5100  
 Houston, Texas 77010  
 (713) 651-5325

*Thomas D Paul*

SIGNATURE: \_\_\_\_\_

Thomas D Paul

NAME \_\_\_\_\_

32,714

REGISTRATION NUMBER \_\_\_\_\_

09/856339

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Serial No.:	To be assigned	§	Docket No.:	P02196US0
		§		
Filed:	May 18, 2001	§		
		§		
Applicant:	Isis Innovation Limited	§	Group Unit:	Not yet assigned
		§		
Title:	Process for Oxidising	§	Examiner:	Not yet assigned
	Terpenes	§		

Box Patent Application  
 Assistant Commissioner for Patents  
 Washington D. C. 20231

## PRELIMINARY AMENDMENT

Dear Sir:

Applicants respectfully request that a Preliminary Amendment for the above-identified Application be entered as contained herein.

In the claims:

In this patent application, please consider under prosecution claims 1-18 wherein amended claims 1, 3, 6, 8-11, and 13-18 are entered as follows:

1. (Amended) Process for oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene or a substituted derivative thereof, and which process comprises:  
 oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain.

3. (Amended) Process according to claim 2 in which the enzyme is one in which amino acid 47 or 51 or 47 and 51 of P450<sub>BM-3</sub>, or amino acid 96 of P450<sub>cam</sub>, or the equivalent amino acid in a said homologue, have been changed to an amino acid with a less polar side-chain.

6. (Amended) The enzyme as defined in claim 4 excluding mutants of P450<sub>cam</sub> which only have the mutations F87A-Y96G-F193A, F87A-Y96G-F193A-C334A, or T101M-T185F-V247M.

8. (Amended) A cell which expresses:

(i) a mutant haem-containing enzyme comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain which in its naturally occurring form has an electron transfer reductase domain, or

(ii) (a) a mutant haem-containing enzyme comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain,

(b) an electron transfer reductase, and

(c) an electron transfer redoxin; or

(iii) (a) (1) P450<sub>cam</sub>, or a fragment thereof; or

(2) a naturally occurring homologue of P450<sub>cam</sub> or a fragment thereof; or

(3) a mutant P450<sub>cam</sub>, or a mutant of a naturally occurring homologue of P450<sub>cam</sub> comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain; or

(4) a P450<sub>cam</sub> which has at least 70% amino acid homology with (1), (2) or (3) and optionally in which amino acid 96 has been changed to an amino acid with a less polar side-chain; and

(b) an electron transfer reductase; and

- (c) an electron transfer redoxin, or
- (iv) (a) (1) P450<sub>BM-3</sub>, or a fragment thereof; or
- (2) a naturally occurring homologue of P450<sub>BM-3</sub> or a fragment thereof; or
- (3) a mutant P450<sub>BM-3</sub>, or a mutant of a naturally occurring homologue of P450<sub>BM-3</sub> comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain

excluding an *E. Coli* DH5 $\alpha$  cell in which the only mutants of P450<sub>cam</sub> which are expressed are amongst the following:

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPGPSST-putidaredoxin reductase-TDGASSS putidaredoxin-COOH,

H<sub>2</sub>N-450<sub>cam</sub>-putidaredoxin reductase-TDGASSS-putidaredoxin-PLEL-P450<sub>cam</sub>-COOH.

9. (Amended) The cell according to claim 8 in which (a), (b) and (c) or (b) and (c) are expressed together in the same fusion protein.

10. (Amended) The cell according to claim 8 in which:

- (b) is putidaredoxin reductase or a fragment thereof; and/or
- (c) is putidaredoxin or a fragment thereof.

11. (Amended ) Process for oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene, or a substituted derivative thereof, and which process comprises oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of a first amino acid in an active site by an amino acid with a less polar side-chain, wherein the substrate is oxidised in a cell according to claim 8.
13. (Amended) Process for selecting a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or a homologue thereof, for its ability to oxidise a substrate, which process comprises screening a group of said mutants for their oxidation effect on the substrate.
14. (Amended) Process according to claim 13 in which the mutant is additionally selected for its ability to oxidise the substrate to an oxidation product.
15. (Amended) Process for selecting a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or a homologue thereof, for its ability to oxidise a substrate, which process comprises screening a group of said mutants for their oxidation effect on the substrate in which the screening is carried out on the library made in a process according to claim 12.
16. (Amended) A process for producing a library of oxidation products comprising providing an acyclic or cyclic terpene, or a cycloalkene, or a substituted derivative thereof to a library made in a process according to claim 12 and allowing oxidation of the substrate.
17. (Amended) A method of treatment of a human or an animal comprising administering an oxidation product obtained by oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene, or a substituted derivative thereof, and which process comprises oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain, wherein optionally the enzyme is one which has been selected in a process according to claim 13.

18. (Amended) A pharmaceutical composition comprising an oxidation product obtained by oxidising a substrate which is an acyclic or a cyclic terpene, or a cycloalkene, or a substituted derivative thereof, and which process comprises oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain, wherein optionally the enzyme is one which has been selected in a process according to claim 13 and a pharmaceutically acceptable carrier or diluent.

**REMARKS**

Applicants have attached a marked version of the claims in this application. For the Examiner's convenience, Applicants have also attached a clean copy of all pending claims in this application after amendments.

Respectfully submitted,



Thomas D. Paul  
Registration No. 32,714  
Counsel for Applicants

Date: 5/18/01  
Fulbright & Jaworski L.L.P.  
1301 McKinney, Suite 5100  
Houston, Texas 77010-3095  
Telephone: 713-651-5325  
Telecopier: 713-651-5246



**MARKED VERSION OF CLAIMS SHOWING AMENDMENTS**

1. (Amended) Process for oxidising a substrate which is an acyclic or cyclic terpene [;] a or a cycloalkene or a substituted derivative thereof, and which process comprises:  
oxidising said [compound] substrate with a mutant haem-containing enzyme, the mutant comprising [the] a substitution of an amino acid in [the] an active site by an amino acid with a less polar side-chain.
3. (Amended) Process according to claim 2 in which the enzyme is one in which amino acid 47 [and/] or 51 or 47 and 51 of P450<sub>BM-3</sub>, or amino acid 96 of P450<sub>cam</sub>, or the equivalent amino acid[(s)] in a said homologue, have been changed to an amino acid with a less polar side-chain
6. (Amended) [An enzyme] The enzyme as defined in claim 4 excluding mutants of P450<sub>cam</sub> which only have the mutations F87A-Y96G-F193A, F87A-Y96G-F193A- C334A, or T101M-T185F-V247M.
8. (Amended) A cell which expresses:
  - (i) a mutant haem-containing enzyme comprising [the] a substitution of an amino acid in [the] an active site by an amino acid with a less polar side-chain which in its naturally occurring form has an electron transfer reductase domain, or
  - (ii)
    - (a) a mutant haem-containing enzyme comprising [the] a substitution of an amino acid in [the] an active site by an amino acid with a less polar side-chain,
    - (b) an electron transfer reductase, and

- (c) an electron transfer redoxin; or
- (iii) (a) (1) P450<sub>cam</sub>, or a fragment thereof; or
  - (2) a naturally occurring homologue of P450<sub>cam</sub> or a fragment thereof; or
  - (3) a mutant P450<sub>cam</sub>, or a mutant of a naturally occurring homologue of P450<sub>cam</sub> comprising [the] a substitution of an amino acid in [the] an active site by an amino acid with a less polar side-chain; or
  - (4) a P450<sub>cam</sub> which has at least 70% amino acid homology with (1), (2) or (3) and optionally in which amino acid 96 has been changed to an amino acid having a less polar side-chain; and
- (b) an electron transfer reductase; and
- (c) an electron transfer redoxin, or
- (iv) (a) (1) P450<sub>BM-3</sub>, or a fragment thereof; or
  - (2) a naturally occurring homologue of P450<sub>BM-3</sub> or a fragment thereof; or
  - (3) a mutant P450<sub>BM-3</sub>, or a mutant of a naturally occurring homologue of P450<sub>BM-3</sub> comprising [the] a substitution of an amino acid in [the] an active site by an amino acid with a less polar side-chain

excluding an *E. Coli* DH5 $\alpha$  cell in which the only mutants of P450<sub>cam</sub> which are expressed are amongst the following:

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPGPSST-putidaredoxin reductase-TDGASSS  
putidaredoxin-COOH,

H<sub>2</sub>N-450<sub>cam</sub>-putidaredoxin reductase-TDGASSS-putidaredoxin-PLEL-P450<sub>cam</sub>-COOH.

9. (Amended) [A] The cell according to claim 8 in which (a), (b) and (c) or (b) and (c) are expressed together in the same fusion protein.
10. (Amended) [A] The cell according to claim 8 in which:
- (b) is putidaredoxin reductase or a fragment thereof; and/or
  - (c) is putidaredoxin or a fragment thereof.
11. (Amended) Process for oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene, or a substituted derivative thereof[:], and which process comprises oxidising said [compound] substrate with a mutant haem-containing enzyme, the mutant comprising [the] a substitution of a first amino acid in [the] an active site by an amino acid with a less polar side-chain, wherein the [compound] substrate is oxidised in a cell according to claim 8.
13. (Amended) Process for selecting a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or a homologue thereof, for its ability to oxidise a [particular] substrate, which process comprises screening a group of said mutants for their oxidation effect on the [particular] substrate.
14. (Amended) Process according to claim 13 in which the mutant is additionally selected for its ability to oxidise the [particular compound] substrate to [a] an [particular] oxidation product.

15. (Amended) Process [according to claim 13] for selecting a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>  
or a homologue thereof, for its ability to oxidise a substrate, which process comprises  
screening a group of said mutants for their oxidation effect on the substrate in which  
the screening is carried out in a library made in a process according to claim 12.
16. (Amended) A process for producing a library of oxidation products comprising  
providing an acyclic or cyclic terpene, or a cycloalkene[;], or a substituted derivative  
thereof to a library made in a process according to claim 12 and allowing oxidation of  
the substrate.
17. (Amended) A method of treatment of a human or an animal comprising administering an  
oxidation product obtained by oxidising a substrate which is an acyclic or cyclic  
terpene, or a cycloalkene[;], or a substituted derivative thereof, and which  
process comprises oxidising said substrate with a mutant haem-containing enzyme,  
the mutant comprising [the] a substitution of an amino acid in [the] an active site by  
an amino acid with a less polar side-chain, wherein optionally the enzyme is one  
which has been selected in a process according to claim 13.
18. (Amended) A pharmaceutical composition comprising an oxidation product obtained by  
oxidising a substrate which is an acyclic or a cyclic terpene, or a  
cycloalkene[;], or a substituted derivative thereof, and which process comprises  
oxidising said [compound] substrate with a mutant haem-containing  
enzyme, the mutant comprising [the] a substitution of an amino acid in [the] an active  
site by an amino acid with a less polar side-chain, wherein optionally the enzyme is  
one which has been selected in a process according to claim 13 and a  
pharmaceutically acceptable carrier or diluent.

CLEAN COPY OF ALL PENDING CLAIMS AS OF MAY 18, 2001

1. (Amended) Process for oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene or a substituted derivative thereof, and which process comprises:  
oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain.
2. Process according to claim 1 in which the enzyme is a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or a mutant of a naturally occurring homologue of either of these enzymes.
3. (Amended) Process according to claim 2 in which the enzyme is one in which amino acid 47 or 51 or 47 and 51 of P450<sub>BM-3</sub>, or amino acid 96 of P450<sub>cam</sub>, or the equivalent amino acid in a said homologue, have been changed to an amino acid with a less polar side-chain.
4. Process according to claim 1 in which there are one or more other amino acid substitutions in the active site.
5. Process according to claim 1 in which the enzyme is (i) P450<sub>cam</sub> and comprises one or more of the following mutations: F87W, F87I, F87L, T185L, T185F, V247A, V247L or F87A-I395F; or (ii) P450<sub>BM-3</sub> and comprises the mutation R47L-Y51F.
6. (Amended) The enzyme as defined in claim 4 excluding mutants of P450<sub>cam</sub> which only have the mutations F87A-Y96G-F193A, F87A-Y96G-F193A-C334A, or T101M-T185F-V247M.
7. A polynucleotide which comprises a sequence which encodes an enzyme as defined in claim 6.

8. (Amended) A cell which expresses:

- (i) a mutant haem-containing enzyme comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain which in its naturally occurring form has an electron transfer reductase domain, or
- (ii)
  - (a) a mutant haem-containing enzyme comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain,
  - (b) an electron transfer reductase, and
  - (c) an electron transfer redoxin; or
- (iii)
  - (a)
    - (1) P450<sub>cam</sub>, or a fragment thereof; or
    - (2) a naturally occurring homologue of P450<sub>cam</sub> or a fragment thereof; or
    - (3) a mutant P450<sub>cam</sub>, or a mutant of a naturally occurring homologue of P450<sub>cam</sub> comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain; or
    - (4) a P450<sub>cam</sub> which has at least 70% amino acid homology with (1), (2) or (3) and optionally in which amino acid 96 has been changed to an amino acid with a less polar side-chain; and
  - (b) an electron transfer reductase; and
  - (c) an electron transfer redoxin, or
- (iv)
  - (a)
    - (1) P450<sub>BM-3</sub>, or a fragment thereof; or
    - (2) a naturally occurring homologue of P450<sub>BM-3</sub> or a fragment thereof; or

(3) a mutant P450<sub>BM-3</sub>, or a mutant of a naturally occurring homologue of P450<sub>BM-3</sub> comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain

excluding an *E. Coli* DH5 $\alpha$  cell in which the only mutants of P450<sub>cam</sub> which are expressed are amongst the following:

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGGPSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGGPSST-putidaredoxin reductase-TDGASSS putidaredoxin-COOH,

H<sub>2</sub>N-450<sub>cam</sub>-putidaredoxin reductase-TDGASSS-putidaredoxin-PLEL-P450<sub>cam</sub>-COOH.

9. (Amended) The cell according to claim 8 in which (a), (b) and (c) or (b) and (c) are expressed together in the same fusion protein.

10. (Amended) The cell according to claim 8 in which:

(b) is putidaredoxin reductase or a fragment thereof; and/or

(c) is putidaredoxin or a fragment thereof.

11. (Amended ) Process for oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene, or a substituted derivative thereof, and which process comprises oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of a first amino acid in an active site by an amino acid with a less polar side-chain, wherein the substrate is oxidised in a cell according to claim 8.

12. A process for making a library of mutants of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or mutants of a homologue of either of these enzymes comprising contacting a cell according to claim 8, or an *E.Coli* DH5 $\alpha$  cell in which the only mutants of P450<sub>cam</sub> which are expressed are amongst the following:

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPGPSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-putidaredoxin reductase-TDGASSS-putidaredoxin-PELEL-P450<sub>cam</sub>-COOH;

with a mutagen and/or when the cell is a mutator cell culturing the cell in conditions in which mutants are produced.

13. (Amended) Process for selecting a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or a homologue thereof, for its ability to oxidise a substrate, which process comprises screening a group of said mutants for their oxidation effect on the substrate.
14. (Amended) Process according to claim 13 in which the mutant is additionally selected for its ability to oxidise the substrate to an oxidation product.
15. (Amended) Process for selecting a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or a homologue thereof, for its ability to oxidise a substrate, which process comprises screening a group of said mutants for their oxidation effect on the substrate in which the screening is carried out on the library made in a process according to claim 12.



16. (Amended) A process for producing a library of oxidation products comprising providing an acyclic or cyclic terpene, or a cycloalkene, or a substituted derivative thereof to a library made in a process according to claim 12 and allowing oxidation of the substrate.
17. (Amended) A method of treatment of a human or an animal comprising administering an oxidation product obtained by oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene, or a substituted derivative thereof, and which process comprises oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain, wherein optionally the enzyme is one which has been selected in a process according to claim 13.
18. (Amended) A pharmaceutical composition comprising an oxidation product obtained by oxidising a substrate which is an acyclic or a cyclic terpene, or a cycloalkene, or a substituted derivative thereof, and which process comprises oxidising said substrate with a mutant haem-containing enzyme, the mutant comprising a substitution of an amino acid in an active site by an amino acid with a less polar side-chain, wherein optionally the enzyme is one which has been selected in a process according to claim 13 and a pharmaceutically acceptable carrier or diluent.

PROCESS FOR OXIDISING TERPENES

The invention relates to a process for enzymatically oxidising terpenes and cycloalkenes.

Terpenoid compounds are widespread in biological systems and constitute one of the largest class of natural products. They are major constituents of essential oils, some of which are of considerable value in the flavour and perfume industries. Many terpenoids are also biologically active. Some are anti-bacterial and anti-fungal agents and thus are of great interest to the pharmaceutical industry. Indeed, terpenoids are some of the highest added value chemicals.

The terpenoids of commercial interest are not normally the terpenes themselves, but rather derivatives which commonly require stereoselective functionalisation at allylic as well as non-activated carbon-hydrogen bonds of the parent terpene. This type of chemical transformation is one of the most difficult reaction to carry out by conventional methods of chemical synthesis - the highly reactive chemical oxidising agents required are non-selective and typically they will preferentially attack more activated carbon-hydrogen bonds and reactive functional groups such as olefinic double bonds commonly present in terpenes.

The present invention concerns the enzymatic oxidation of terpenes and cycloalkenes. This technique enables the synthesis of hydroxylated terpenes (and cycloalkenes), often in a single step, and provided that the match between substrate and enzyme is correct, the oxidation reaction can be highly chemoselective (attack at a particular functional group such as a non-activated C-H bond rather than some other reactive functional group) and stereoselective. The fine tuning and alterations of substrate specificity and selectivity of substrate oxidation are very difficult to achieve for

-2-

conventional reagents.

The present invention provides a process for oxidising a substrate which is an acyclic or cyclic terpene or a cycloalkene, or a substituted derivative thereof, which process comprises oxidising said compound with a mutant haem-containing enzyme, the mutant comprising the substitution of an amino acid in the active site by an amino acid with a less polar side-chain.

Although the terpenes used in the present invention will generally have the formula  $(C_5H_8)_n$  where  $n$  is 2 or more, especially 2, 3 or 4, it is to be understood that the term "terpene" extends to compounds which are strictly referred to as "terpenoid", involving the loss or shift of a fragment, generally a methyl group. Thus, for example, sesquiterpenes (where  $n$  is 3) which can be used in the present invention may contain only, say, 14, rather than 15, carbon atoms. Generally the terpene is one which can be built up from isoprene units. The terpene may be cyclic or acyclic.

The monoterpenes (where  $n$  is 2) will generally have 10 carbon atoms, typically with 1 to 3 double bonds, especially 1 or 2 ring double bonds, and typically with 0 to 2 rings. It is possible for one of the rings to be formed as a bridge containing, typically 0 or 1 carbon atoms. In other words, it can be formed by a direct link between 2 carbon atoms of an existing ring or with an intermediate methylene group. If the terpene is acyclic it will generally contain at least 2 double bonds and generally 3.

The sesquiterpenes will normally contain 14 or 15 carbon atoms, typically with 0 to 2 double bonds and typically 1 to 3 rings, with the possibility of fused rings and/or bridged rings.

The rings which may be present in the terpenes will

-3-

typically have from 3 to 9 carbon atoms, more especially 5 or 6 carbon atoms. Thus, in particular, the terpenes will contain a cyclohexane, or cyclohexadiene ring.

5 The terpenes will generally contain a total of 3 or 4 exocyclic methyl or methylene groups, for example 2 methyl groups and 1 methylene group or 3 methyl groups for a monoterpene, and 3 methyl groups and 1 methylene group or 4 methyl groups for a sesquiterpene.

10 The monoterpene is typically a limonene, pinene, terpinene, sabinene, thujene, mercene, ocimene, nerol or geraniol, for example as shown in Table 1.

15 The sesquiterpene is generally formed by a head-to-tail arrangement of three isoprene units. The sesquiterpene is typically an aromadendrene, caryophyllene, longifolene, valencene, isobazzanene, silphinene, ishwarane, isopatchchoul-3-ene, or isosesquicarene, for example as shown in Table 2.

The diterpene (where n is 4) is typically casbene, retinal, abietic acid or a gibberellin.

20 The cycloalkene generally comprises up to 9 ring members, e.g. it is a 5, 6, 7, 8, 9 or more membered ring. The cycloalkene is typically a cyclohexene.

25 Substituted derivatives of any of the terpenes or cycloalkenes mentioned above may also be used. Typically 1, 2, 3 or more substituents are present. Any combination of the following substituents may be present. The substituent is typically a halogen atom or an alkyl or alkenyl group, which generally has 1 to 6 carbons, the substituent optionally being substituted with one or more  
30 halogens. It is generally not phenylcyclohexene. Indeed the presence of aromatic, such as phenyl, rings is generally avoided for all the substrates used in the invention.

35 The substituent typically has the formula  $C_nH_kX_m$ , wherein X is the halogen, n is 1, 2, 3 or more, m is 1,

-4-

2, 3, 4 or more and  $k$  is an integer which has an appropriate value so that the valencies of the substituent  $C_nH_kX_m$  are satisfied. For an alkyl substituent  $k+m = 2n+1$ . Typically  $k$  is 1, 2, 3, 4 or more, or may be 0, i.e. the substituent is a perhaloalkyl group. The halogen is typically fluorine, chlorine or bromine.

The substituent may also comprise 1, 2 or more oxygen atoms and for example may be an alcohol, aldehyde, ketone or epoxide group.

The oxidation causes the formation of a C-O bond in the compound, generally as the hydroxide from the oxidation of a carbon-hydrogen bond, but an epoxide may be formed from the oxidation of a C=C bond. The oxidation may thus introduce a hydroxy, aldehyde, ketone or epoxide group. Alternatively the oxidation may cause the further oxidation of an oxygen containing group, such as converting a hydroxy group into an aldehyde or ketone group. 1, 2 or more carbon atoms may be attacked in the same substrate molecule.

The oxidation typically gives rise to 1, 2 or more oxidation products. These different products may result from different carbon atoms being attacked and/or from different degrees of oxidation occurring at a given carbon atom.

The oxidation may occur on either a ring carbon atom or a substituent carbon atom or both. At least the initial oxidation will involve attack of a C-H bond which may be activated or non-activated or attack at a carbon-carbon double bond (typically giving an epoxide). Generally an activated C-H bond is where the carbon atom is in a benzylic or allyl position. Aromatic rings and olefinic double bonds activate C-H bonds to attack by stabilising the radical intermediate or any build-up of charge generated during the reaction pathway. The

-5-

carbon of the C-H bond may be a primary, secondary or tertiary carbon.

The oxidation typically preserves stereoisomerism. Thus when the substrate consists of a single stereoisomer the product typically consists of a single corresponding stereoisomer, or can contain a preponderance of the corresponding stereoisomer.

The enzyme used in the process is generally a P450 enzyme, typically of eukaryotic or prokaryotic origin. The enzyme is generally of bacterial, fungal, yeast, plant or animal origin, and thus may be from a bacterium of the genus *Pseudomonas*. The enzyme is typically a monooxygenase. The non-mutant form of the enzyme may or may not be able to oxidize terpenes and/or cycloalkenes.

The mutations discussed herein are generally introduced into the enzyme by using methods known in the art, such as site directed mutagenesis of the enzyme, PCR and gene shuffling methods or by the use of multiple mutagenic oligonucleotides in cycles of site-directed mutagenesis. Thus the mutations may be introduced in a directed or random manner. The mutagenesis method thus produces one or more polynucleotides encoding one or more different mutants. Typically a library of mutant oligonucleotides is produced which can be used to produce a library of mutant enzymes.

An amino acid 'in the active site' is one which lines or defines the site in which the substrate is bound during catalysis or one which lines or defines a site through which the substrate must pass before reaching the catalytic site. Therefore such an amino acid typically interacts with the substrate during entry to the catalytic site or during catalysis. Such an interaction typically occurs through an electrostatic interaction (between charged or polar groups), hydrophobic interaction, hydrogen bonding or van der Waals forces.

-6-

The amino acids in the active site can be identified by routine methods to those skilled in the art. These methods include labelling studies in which the enzyme is allowed to bind a substrate which modifies ('labels') amino acids which contact the substrate. Alternatively the crystal structure of the enzyme with bound substrate can be obtained in order to deduce the amino acids in the active site.

The enzyme may have 1, 2, 3, 4, 5 to 10, 10 to 20 or more other mutations, such as substitutions, insertions or deletions. The other mutations may be in the active site or outside the active site. Typically the mutations are in the 'second sphere' residues which affect or contact the position or orientation of one or more of the amino acids in the active site. The insertion is typically at the N and/or C terminal and thus the enzyme may be part of a fusion protein. The deletion typically comprises the deletion of amino acids which are not involved in catalysis, such as those outside the active site (thus the enzyme is a mutated fragment of a naturally occurring enzyme). The enzyme may thus comprise only those amino acids which are required for oxidation activity.

The other mutation in the active site typically alters the position and/or conformation of the substrate when it is bound in the active site. The mutation may make the site on the substrate which is to be oxidized more accessible to the haem group. Thus the mutation may be a substitution to an amino acid which has a smaller or larger, or more or less polar, side chain.

The other mutations typically increase the stability of the protein, or make it easier to purify the protein. They typically prevent the dimerisation of the protein, typically by removing cysteine residues from the protein (e.g. by substitution of cysteine at position 334

-7-

of P450<sub>cam</sub>, or at an equivalent position in a homologue, preferably to alanine). They typically allow the protein to be prepared in soluble form, for example by the introduction of deletions or a poly-histidine tag, or by mutation of the N-terminal membrane anchoring sequence. The mutations typically inhibit protein oligomerisation, such as oligomerisation arising from contacts between hydrophobic patches on protein surfaces.

Thus the mutant enzyme is typically at least 70% homologous to a naturally occurring haem-containing enzyme on the basis of amino acid identity.

Any of the homologous proteins (i.e. described as being homologous to another protein) mentioned herein are typically at least 70% homologous to the relevant protein or at least 80 or 90% and more preferably at least 95%, 97% or 99% homologous thereto over at least 20, preferably at least 30, for instance at least 40, 60 or 100 or more contiguous amino acids. The contiguous amino acids may include the active site. This homology may alternatively be measured not over contiguous amino acids but over only the amino acids in the active site.

Homology can be measured using known methods. For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology (for example used on its default settings) (Devereux et al (1984) *Nucleic Acids Research* 12, p387-395). The PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (typically on their default settings), for example as described in Altschul S. F. (1993) *J Mol Evol* 36:290-300; Altschul, S, F et al (1990) *J Mol Biol* 215:403-10.

Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring



-8-

sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul et al, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

-9-

Typically the homologous protein differs from the relevant protein by at least 1, 2, 5 or 10 mutations (substitutions, insertions or deletions) when compared to all of the protein or over any of the lengths of contiguous amino acids mentioned above.

The enzyme used in the process is preferably a mutant of P450<sub>cam</sub> (such as mutant of the sequence shown in table 7) or a mutant of a naturally occurring homologue of P450<sub>cam</sub>, typically of P450<sub>BM-3</sub> from *Bacillus megaterium* (such as a mutant of the sequence shown in table 8), P450<sub>terp</sub> from *Pseudomonas sp.*, and P450<sub>eryF</sub> from *Saccharopolyspora erythraea*, and also P450 105 D1 (CYP105) from *Streptomyces griseus* strains. Note that the amino acid numbering shown in table 8 for P450<sub>BM-3</sub> does not correspond to the numbering used in the description to denote mutations in this enzyme. The sequence shown in table 8 contains an additional amino acid at the N terminal. This is normally cleaved in vivo. Therefore each amino acid number shown in the table is always one more than the number used in the conventional numbering (as used in the description).

The naturally occurring homologue of P450<sub>cam</sub> (e.g. of P450<sub>BM-3</sub>) may have substantially the same activity as P450<sub>cam</sub> or P450<sub>BM-3</sub>. The homologue may be a species homologue or an allelic variant of P450<sub>cam</sub> from *Pseudomonas putida* or of P450<sub>BM-3</sub>. The amino acids in the active site of the homologue may be the same as in the active site of P450<sub>cam</sub> or of P450<sub>BM-3</sub>. Typically the amino acid at the equivalent position to 96 in P450<sub>cam</sub> is a tyrosine in the homologue.

The mutant of P450<sub>cam</sub> or of a homologue of P450<sub>cam</sub> is typically one in which amino acid 96, or the equivalent amino acid in a homologue, has been changed to an amino acid with a less polar side chain. In the case where the homologue is P450<sub>BM-3</sub> the mutant typically has a

-10-

substitution (to a less polar amino acid) at 47 and/or 51 and/or 42 and/or 75 and/or 354 and/or 264 and/or 263 and/or 181 and typically does not have a mutation at the equivalent site to 96 of P450<sub>cam</sub> (preferred mutants of P450<sub>BM-3</sub> have at least mutations at 47 and 51, or at the equivalent sites in homologues).

Thus typically the substitution is to an amino acid which is above the original amino acid in Table 3, such as the preferred mutations shown in Table 4 and Table 5.

The 'equivalent' side chain in the homologue is one at the homologous position. This can be deduced by lining up the P450<sub>cam</sub> or P450<sub>BM-3</sub> sequence and the sequence of the homologue based on the homology between the two sequences. The PILEUP and BLAST algorithms can be used to line up the sequences. The equivalent amino acid will generally be in a similar place in the active site of the homologue as any of the specific amino acids discussed herein, such as amino acid 96 in P450<sub>cam</sub>.

The discussion below provides examples of the positions at which substitutions may be made in P450<sub>cam</sub> and P450<sub>BM-3</sub>. The same substitutions may be made at equivalent positions in the homologues. Standard nomenclature is used to denote the mutations. The letter of the amino acid present in the natural form is followed by the position, followed by the amino acid in the mutant (these positions can be correlated to the numbering shown in tables 7 and 8 with the proviso discussed above with regard to table 8 amino acid numbering). To denote multiple mutations in the same protein each mutation is listed separated by hyphens. The mutations discussed below using this nomenclature specify the natural amino acid in P450<sub>cam</sub> or P450<sub>BM-3</sub> but it is to be understood that the (same) mutation could be made to a homologue which has a different amino acid at the equivalent position.

An additional mutation is typically an amino acid

-11-

substitution at amino acid 87, 98, 101, 185, 244, 247, 248, 296, 395, 396 of P450<sub>cam</sub> (or a combination of these, for example as shown in table 4).

The following combinations of substitutions are preferred for P450<sub>cam</sub>:

(i) Substitution at position 87 to amino acids of different side-chain volume, such as substitutions (typically of F) to A, L, I and W, combined with substitutions at position 96 to amino acids of different side-chain volume such as (typically Y to) A, L, F, and W. These combinations alter the space available in the upper part of the substrate pocket compared to the wild-type enzyme, for example, from Y96W-F87W (little space) to Y96A-F87A (more space), as well as the location of the space, for example from one side in Y96F-F87A to the other in Y96A-F87W.

(ii) Substitution at position 96 to F combined with substitutions at positions 185 and 395. Both T185 and I395 are at the upper part of the substrate pocket, and substitution with A creates more space while substitution with F will reduce the space available and push the substrate close to the haem.

(iii) Substitutions at position 96 to A, L, F, and W combined with substitutions at residues closer to the haem including at 101, 244, 247, 295, 296 and 396 to A, L, F, or W. These combinations will create or reduce space in the region of the different side-chains to offer different binding orientations to substrates of different sizes. For example, the combinations Y96W-L244A and Y96L-V247W will offer very different substrate pockets for the binding of R-limonene.

(iv) Triple substitutions at combinations of positions 87, 96, 244, 247, 295, 296, 395 and 396 with combinations of A, L, F, and W. The aim is to vary the size and shape of the hydrophobic substrate binding pocket. For

-12-

example, the Y96A-F87A-L244A combination creates more space compared to the Y96F-F87W-V396L combination, thus allowing larger terpenes to bind to the former while restricting the available binding orientations of smaller terpenes in the latter. The combinations Y96F-F87W-V247L and Y96F-F87W-V295I have comparable substrate pocket volumes, but the locations of the space available for substrate binding are very different. The combination Y96F-F87L-V247A has a slightly larger side-chain volume at the 96 position than the combination Y96L-F87L-V247A, but the L side-chain at the 96 position is much more flexible and the substrate binding orientations will be different for the two triple mutants.

(v) The mutants with four or five substitutions were designed with similar principles of manipulating the substrate volume, the different flexibility of various side-chains, and the location of the space available in the substrate pocket for terpene binding so as to effect changes in selectivity of substrate oxidation.

The invention also provides the mutant of P450<sub>cam</sub> or a mutant of a homologue of P450<sub>cam</sub> (such as P450<sub>BM-3</sub>) as discussed above, excluding mutants of P450<sub>cam</sub> which only have the mutations:

F87A-Y96G-F193A, F87A-Y96G-F193A-C334A or T101M-T185F-V247M.

The mutant enzyme may be in a substantially isolated form and/or a substantially purified form, in which case it will generally comprise (e.g. about or at least) 90%, such as (e.g. about or at least) 95%, 98% or 99% of the protein in the preparation.

The invention also provides a polynucleotide which comprises a sequence which encodes the mutant enzyme of the invention. The polynucleotide is typically DNA or RNA, and may be single or double stranded. The polynucleotide may be able to hybridise with a

-13-

polynucleotide encoding the naturally occurring form of any mutant discussed herein (each the polynucleotide shown in table 7 or 8). It typically hybridises with the relevant polynucleotide at a level significantly above background. The signal level generated by the interaction is typically at least 10 fold, preferably at least 100 fold, as intense as 'background' hybridisation. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with  $^{32}\text{P}$ . Selective hybridisation is typically achieved using conditions of medium to high stringency (for example 0.03M sodium chloride and 0.03M sodium citrate at from about 50°C to about 60°C).

Polynucleotides of the invention can be incorporated into a recombinant replicable vector. The vector may be used to replicate the nucleic acid in a compatible host cell. Thus one method of making polynucleotides of the invention comprises introducing a polynucleotide of the invention into a replicable vector, introducing the vector into a compatible host cell, and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell. Suitable host cells are described below in connection with expression vectors.

Preferably, a polynucleotide of the invention in a vector is operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector.

The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under

-14-

conditions compatible with the control sequences.

Such vectors may be transformed into a suitable host cell to provide for expression of the mutant enzyme.

The vectors may be, for example, plasmid, virus or phage vectors provided with an origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a neomycin resistance gene for a mammalian vector. Vectors may be used *in vitro*, for example for the production of RNA or used to transfect or transform a host cell.

Promoters and other expression regulation signals may be selected to be compatible with the host cell for which the expression vector is designed. For example, *E. Coli* promoters include *lac*, *tac*, *trc*, *trp* and T7 promoters, and yeast promoters include *S. cerevisiae* GAL4 and ADH promoters, *S. pombe* *nmt1* and *adh* promoters. Mammalian promoters include the metallothionein promoter which can be induced in response to heavy metals such as cadmium. The expression vectors are possible for use in insect or mammalian cells. For use in insect cells, strong baculovirus promoters such as the polyhedrin promoter are preferred. For expression in mammalian cells, strong viral promoters such as the SV40 large T antigen promoter, a CMV promoter or an adenovirus promoter may also be used. All these promoters are readily available in the art.

Expression vectors of the invention are typically introduced into host cells using conventional techniques including calcium phosphate precipitation, DEAE-dextran transfection, or electroporation.

The expression vector may contain a selectable marker and/or such a selectable marker may be co-

-15-

transfected with the expression vector and stable transfected cells may be selected.

Suitable cells include cells in which the abovementioned vectors may be expressed. Such cells may be prokaryotic or eukaryotic. These include microbial cells typically bacteria such as *E. coli*, preferably the strains DH<sub>5α</sub>, JM109, NM522 and BL21DE3 or *Pseudomonas*, typically putida, mammalian cells such as CHO cells, COS7 cells or HeLa cells, insect cells or yeast such as *Saccharomyces*. Baculovirus or vaccinia expression systems may be used.

Cell culture can take place under standard conditions. Generally the cells are cultured in the presence of assimilable carbon and nitrogen sources. Commercially available culture media for cell culture are widely available and can be used in accordance with manufacturers instructions.

Typically the process of the invention is carried out in vitro, such as in a cell free system. The process may be carried out in vivo in a cell.

Typically, in addition to the enzyme (a) and the substrate the process of the invention is carried out in the presence of an electron transfer reductase (b), an electron transfer redoxin (c), cofactor for the enzyme and an oxygen donor. In this system the flow of electrons is typically: cofactor → (b) → (c) → (a). However particular enzymes do not require the presence of an electron reductase and electron transfer redoxin, such as P450<sub>BM-3</sub>. Although the following discussion is particularly directed to enzymes which do require a reductase or redoxin, it is applicable to enzymes which do not require these (for example the various concentrations, conditions and rates are suitable for these enzymes).

For enzymes which do require a reductase and



-16-

redoxin (b) is generally an electron transfer reductase which is able to mediate the transfer of electrons from the cofactor to (c), such as a naturally occurring reductase or a protein which has homology with a naturally occurring reductase, such as at least 70% homology; or a fragment of the reductase or homologue. Thus (b) may be derived from any of the organisms listed above from which the haem-containing enzyme may be derived. (b) is typically a flavin dependent reductase, such as putidaredoxin reductase.

(c) is generally an electron transfer redoxin which is able to mediate the transfer of electrons from the cofactor to (a) via (b). (c) is typically a naturally occurring electron transfer redoxin or a protein which has homology with a naturally occurring electron transfer redoxin, such as at least 70% homology; or a fragment of the redoxin or homologue. Thus (c) may be derived from any of the organisms listed above from which the haem-containing enzyme may be derived. (c) is typically a two-iron/two sulphur redoxin, such as putidaredoxin.

The cofactor is any compound capable of donating an electron to (b), such as NADH. The oxygen donor is any compound capable of donating oxygen to (a), such as dioxygen.

In the process the concentration of (a), (b) or (c) is typically from  $10^{-8}$  to  $10^{-2}$ M, preferably from  $10^{-6}$  to  $10^{-4}$ M. Typically the ratio of concentrations of (a): (b) and/or (a): (c) is from 0.1:0.1 to 1:10, preferably from 1:0.5 to 1:2, or from 1:0.8 to 1:1.2. Generally the process is carried out at a temperature and/or pH at which the enzyme is functional, such as when the enzyme has at least 20%, 50%, 80% or more of peak activity. Typically the pH is from 3 to 11, such as 5 to 9 or 6 to 8, preferably 7 to 7.8 or 7.4. Typically the temperature is 15 to 90°C, such as 25 to 75°C or 30 to 60°C. In one

-18-

following:

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPGPSST-putidaredoxin reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-putidaredoxin reductases-TDGASSS-putidaredoxin-PLEL-P450<sub>cam</sub>-COOH.

However it is understood that the excluded *E.Coli* DH5α cells can be used to produce the libraries discussed below.

A preferred cell (second type) is a cell which expresses:

- (a) (i) P450<sub>BM-3</sub>, or a fragment thereof; or
- (ii) a naturally occurring homologue of P450<sub>BM-3</sub> or a fragment thereof; or
- (iii) a mutant P450<sub>BM-3</sub>, or a mutant homologue of thereof.

The cell provided by the invention is typically a cell from the species mentioned above in which the nucleotide of the invention can be expressed. The cell may be a mutator cell. Such a cell is generally deficient in one or more of the primary DNA repair pathways (such as *E.Coli* pathways mutS, mutD or mutT, or their equivalents in another organism), and thus has a high mutation rate. Simply culturing such cell leads to the DNA encoding (a) to become mutated. The cell may be of *E.Coli* XL1 Red mutator strain.

The cell of the invention may be in a substantially isolated form and/or substantially purified form, in which case it will generally comprise (e.g. at least or about) 90%, such as (e.g. at least or about) 95%, 98% or 99% of the cells or the mass (normally measured in terms of dry mass) of the preparation.

-17-

embodiment the process is carried out in the presence of a substance able to remove hydrogen peroxide by-product (e.g. a catalase).

Alternatively the process of the invention could be carried out in the presence of the enzyme, substrate and an oxygen atom donor, such as hydrogen peroxide or t-butylhydroperoxide. Thus, the process could be carried out using the peroxide shunt.

Typically in the process at least 20 turnovers/min occur, such as at least 50, 100, 200, 300, 500 or more turnovers (turnover is measured as nanomoles of product formed per nanomole of enzyme).

The invention also provides several types of cells. The first type expresses :

an enzyme which can be used in the process which in its naturally occurring form has an electron transfer reductase domain; or expresses

- (a) the mutant haem-containing enzyme which is used in the process of the invention;
- (b) an electron transfer reductase; and
- (c) an electron transfer redoxin.

The second type of cell expresses:

- (a) (i) P450<sub>cam</sub> or a fragment thereof; or
- (ii) a naturally occurring homologue of P450<sub>cam</sub> or a fragment thereof; or
- (iii) a mutant of P450<sub>cam</sub>; or
- (iii) a polypeptide which has at least 70% amino acid homology with (i) or (ii) and optionally has any of the combination of mutations discussed herein; and
- (b) an electron transfer reductase; and
- (c) an electron transfer redoxin:

excluding an *E.Coli* DH5 $\alpha$  cell in which the only mutants of P450<sub>cam</sub> which are expressed are amongst the

-19-

The cell may be one which does not naturally express (a), (b) or (c). The cell be one in which (a), (b) or (c) are expressed at a higher level than in the naturally occurring cell. (a) may originate from the same organism as (b) or (c).

In the cell (a), (b) and (c) may be expressed from the same vector, or may be expressed from different vectors. They may be expressed as three different polypeptides. Alternatively they may be expressed in the form of fusion proteins. Typically components (a), (b) and (c) are all present in the same fusion protein. Alternatively only two of the components, preferably (b) and (c), may be present in the fusion protein. Typically the components are contiguous in the fusion protein and there is no linker peptide present.

Alternatively a linker may be present between the components. The linker generally comprises amino acids that do not have bulky side chains and therefore do not obstruct the folding of the protein subunits. Preferably the amino acids in the linker are uncharged. Preferred amino acids in the linker are glycine, serine, alanine or threonine. In one embodiment the linker comprises the sequence N-Thr-Asp-Gly-Gly-Ser-Ser-Ser-C. The linker is typically from at least 5 amino acids long, such as at least 10, 30 or 50 or more amino acids long.

The first type of cell may be obtained by transforming or transfecting a host cell with a polynucleotide or vector of the invention.

The mutant enzyme of the invention may be prepared by a process comprising cultivating the first type of cell under conditions to provide for expression of the said mutant enzyme, and optionally recovering the expressed mutant enzyme.

The process of the invention may be carried out in the first type of cell of the invention or in the second

-20-

type of cell if it is able to oxidize the substrate, or a medium containing it. Generally such a process comprises providing the substrate in the cell, allowing the substrate to be oxidized in the process of the invention, and optionally obtaining the oxidation product therefrom, e.g. by extraction. The substrate is typically provided in the cell by adding the substrate to the outside of the cell and allowing it to enter the cell. Alternatively the substrate could be synthesized in the cell from a precursor.

The invention also provides a process for making a library of mutants of P450<sub>cam</sub> or mutants of a homologue of P450<sub>cam</sub> comprising contacting the second type of cell with a mutagen and/or when the cell is a mutator cell culturing the cell in conditions in which mutants are produced. The mutagen may be contacted with the cell prior to or during culturing of the cell. Thus the mutagen may be present during replication of the cell or replication of the genome of the cell.

The mutagen generally causes random mutations in the polynucleotide sequence which encodes (a). The mutagen is typically a chemical mutagen, such as nitrosomethylguanidine, methyl- or ethylmethane sulphonic acid, nitrite, hydroxylamine, DNA base analogues, and acridine dyes, such as proflavin. It is typically electromagnetic radiation, such as ultra-violet radiation at 260 nm (absorption maximum of DNA) and X-rays. It is typically ionising radiation.

Typically the library will be in the form of cells which are derived from cells of the invention by mutagenesis and which cells comprise the mutant enzymes. Generally each cell will express only one particular mutant enzyme. The library typically comprises at least 500 mutants, such as at least 1,000 or 5,000 mutants, preferably at least 10,000 different mutants.

-21-

The library typically comprises a random population of mutants. The library may undergo one or more rounds of selection whilst being produced and therefore may not comprise a random population. Between rounds of selection the cells in the library may be allowed to replicate, and they may also be contacted with a mutagen.

A mutant can be selected from the library based on a particular property of the mutant. The property may comprise one or more of the following characteristics:

- (i) the ability to oxidize a particular substrate; optionally to a particular oxidation product or to a product with a particular activity.
- (ii) the ability to carry out the oxidation of substrate at an increased rate,
- (iii) a reduced oxidation activity towards a particular substrate,
- (iv) a reduction in the production of a particular substrate.

Typically the activity of the product in (i) is blocking the action of an agent which is lethal to the cells of the library. This can be selected by growing the library in the presence of the agent. The agent is typically expressed within the cells of the library.

The activity may be the binding of the product to a particular substance, such a protein. The substance is typically present in the cell of the library, and/or is typically a disease-causing or therapeutic target. An indicator which binds the substance is typically used to detect binding of the product to the substance. In one embodiment the indicator is able to bind the substance and has a property which changes upon binding, e.g. a colour change. Product which displaces the indicator from the substance can thus be detected by measuring changed in the property.

The invention also provides a method of making a

-22-

library of oxidation products comprising providing a substrate to the library of mutant enzymes and allowing oxidation of the substrate.

Products produced in the process of the invention, or identified, selected, made or designed using the library could be used in therapy or diagnosis. Thus the invention provides a method of treating a host suffering from a disease, which method comprises administering to the host a therapeutically effective amount of the product. The condition of a patient suffering from the disease and in need of the product can therefore be improved by administration of the product. The product can also be given as a prophylactic, typically to a host which is at risk from or susceptible to the disease.

The invention provides the product for use in a method of treatment of the human or animal body by therapy. The invention also provides the product for use in a diagnostic method practiced on the human or animal body. The invention also provides use of the product in the manufacture of a medicament to treat a disease.

The formulation of the product for use in preventing or treating infection by an organism will depend upon factors such as the nature of the product identified, whether a pharmaceutical or veterinary use is intended, etc. In order to be administered to a patient, the compound will be provided in the form of a pharmaceutical composition containing the product and a pharmaceutically acceptable carrier or diluent. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. Typical oral dosage compositions include tablets, capsules, liquid solutions and liquid suspensions. For example it may be formulated for parenteral, intravenous, intramuscular, subcutaneous, transdermal or oral administration.

The dose of product may be determined according to

-23-

various parameters, especially according to the substance used; the age, weight and condition of the patient to be treated; the route of administration; and the required regimen. A physician will be able to determine the required route of administration and dosage for any particular patient. A suitable dose may however be from 0.1 to 100 mg/kg body weight such as 1 to 40 mg/kg body weight, for example, to be taken from 1 to 3 times daily.

The invention is illustrated by the accompanying drawings which show the gas chromatography results for various oxidation reactions. In the drawings unless stated otherwise the y-axis shows mVolts and the x-axis shows Time (in minutes).

In the drawings:

Figure 1 shows the oxidation of camphor by the C334A mutant of P450<sub>cam</sub> (expressed from plasmid SGB++). Lines A, B, C, D and E represent camphor turnover at 2, 10, 20, 40 and 100 minutes. The 5.28 peaks is camphor, 11.82 is 5-exo-hydroxycamphor, 7.45 is 5-ketocamphor and the 16.07 peak is the internal standard.

Figure 2 shows R- and S- limonene with wild type P450<sub>BM-3</sub>.

Figure 3 shows NADH consumption by wild type and mutant P450<sub>BM-3</sub>.

Figure 4 shows the whole cell *E. coli* oxidation of  $\alpha$ -pinene by the Y96F-F87W-V247L mutant expressed by plasmid pCWSGB+.

The invention is also illustrated by the Examples:

#### Example 1

##### Expression of mutants for in vitro work.

The P450<sub>cam</sub> enzymes were expressed using the vector pRH1091 (Baldwin, J.E., Blackburn, J.M., Heath, R.J., and Sutherland, J.D. *Bioorg. Med. Chem. Letts.*, 1992, 2, 663-668.) which utilised the *trc* promoter (a fusion of the *trp* and *lac* promoters). This vector incorporates a strong



-24-

ribosome binding site (RBS) and the gene to be expressed is cloned using an *Nde* I site on the 5' end of the gene. We used *Hind* III as the cloning site at the 3' end of the *camC* gene. The procedure for protein expression is as follows: Cells are grown at 30°C until the OD<sub>600nm</sub> reaches 1.0 - 1.2, the temperature is increased to 37°C and camphor added as a 1 M stock in ethanol to a final concentration of 1 mM. The culture is allowed to incubate at 37°C for another 6 hours. The P450<sub>cam</sub> protein is expressed to high levels in the cytoplasm and the cells take on a red to orange-red colour.

We have also prepared a variant of pRH1091 (by PCR) which has a extra *Xba* I site between the RBS and the *Nde* I site. This is important because *Nde* I is not unique in M13, and this restriction site is also present in the reductase gene as well as the backbone of the pGLW11 vector used for the *in vivo* system. *Xba* I is unique in the polylinker region of M13, but absent in the genes of all three proteins in the P450<sub>cam</sub> system and in the expression vectors. It therefore allows the *camC* gene to be moved between the mutagenic and expression vectors.

The P450<sub>BM-3</sub> enzyme from *Bacillus megaterium* was expressed using either the pGLW11 or pCW vectors. The recombinant plasmid with the P450<sub>BM-3</sub> gene inserted into either of these vectors were transformed into *E. coli* strain DH5α and grown under ampicillin selection. A single colony was then grown at 30°C in LB media supplemented with ampicillin until the OD at 600 nm reached ca. 1, and protein expression was induced by adding IPTG from a 1 M stock to a final concentration of 1mM. After 6-8 h, cells were harvested by centrifugation and the expression levels were high, as indicated by an orange-red coloration of the cells.

-25-

How the mutants were made.

Oligonucleotide-directed site-specific mutagenesis was carried out by the Kunkel method (Kunkel, T. A. *Proc. Natl. Acad. Sci. USA* 1985, **82**, 488-492) using the Bio-Rad Mutagen kit and by PCR using the QuikChange kit from Stratagene. The recommended procedure for the Kunkel method is summarised as follows. An M13 mp19 subclone of the *camC* gene encoding P450<sub>cam</sub> was propagated in the *E. coli* strain CJ236. This strain has the *dut*<sup>-</sup>*ung*<sup>-</sup> phenotype and thus will tolerate the inclusion of uracil in place of thymine in DNA molecules. After three cycles of infection, uracil-containing single stranded (USS) M13 DNA was readily isolated by phenol extraction of mature M13 phage particles excreted into the growth medium. The mutagenic oligonucleotide (or oligonucleotides) were phosphorylated with T4 polynucleotide kinase and then annealed to the USS template. The four nucleotides, DNA polymerase, DNA ligase, ATP and other chemical components were added and the second strand was synthesised *in vitro*. The double stranded form thus obtained was transformed into the *dut*<sup>+</sup> *ung*<sup>+</sup> *E. coli* strain MV1190, which should degrade the uracil-containing template strand and propagate the mutant strand synthesised *in vitro*. Plaques were picked and phages of possible mutants grown in *E. coli* strains MV1190 or TG1. The single-stranded DNA from these were sequenced to determine whether the mutagenesis reaction was successful. The mutagenic efficiency was 50 - 80%. The mutant *camC* gene is excised from the M13 subclone by restriction digest with *Nde* I and *Hind* III, and the fragment of appropriate size is ligated to the backbone of the expression vector prepared by a similar *Nde* I/*Hind* III digest.

The QuikChange kit relies on the property of the *Dpn* I restriction enzyme which selectively cleaves methylated

-26-

DNA. The mutation is introduced by PCT using double stranded plasmid DNA, and hence no single stranded template preparations are necessary. The PCR reaction is carried out with two oligonucleotides, one of which binds to the coding strand and the other to the sense strand. Each oligonucleotide contains a short stretch of polynucleotide complementary to either side of the mutation site. After *in vitro* synthesis by PCR using non-methylated dNTP's, plasmid DNA with overlapping nicks in each strand were digested with *Dpn* I to remove the starting template selectively - plasmid DNA isolated from most *E. coli* strains contain methylated bases but the newly synthesised DNA do not have methylated bases. After the digest the DNA is transformed into supercompetent *E. coli* XL1 Blue cells and propagated. The plasmid DNA from potential mutants which grow on agar plates under antibiotic selection were isolated and sequenced to confirm mutagenesis. The cells can then be used for protein expression once the entire sequence of the new mutant was confirmed to ensure that there were no spurious mutations.

Multiple mutants were prepared either by further mutagenesis, also by the Kunkel method, or where the location of the sites in the sequence permits, simple cloning steps. There are two unique restriction sites within the *camC* gene which are absent from the expression vector. One is *Sph* I which spans residues 121 - 123, and the other is *Sal* I which spans residues 338 and 339. Therefore, all mutations at, for example, residues 87, 96, 98, and 101 are readily combined with mutations at higher number residues by ligating appropriate fragments from restriction digests of mutant *camC* genes with *Nde* I/*Sph* I and *Sph* I/*Hind* III and the backbone fragment from a *Nde* I/*Sph* I digest of the expression vector. Mutations

-27-

at, for example, 395 and 396 can be similarly incorporated by digests in which *Sph* I is replaced with *Sal* I.

The rationale for introducing the unique *Xba* I site is now clear: many mutants with multiple mutations were prepared by the cloning procedure above. Without the *Xba* I site it would be impossible to clone the gene for these multiple mutants from the expression vector back into M13 for further rounds of mutagenesis. Of course these problems could be overcome by doing mutagenesis by PCR, for example.

#### Example 2

##### Substrate oxidation protocol: in vitro reactions

Component	Final concentration
P450 <sub>cam</sub> enzyme	1 $\mu$ M
Putidaredoxin	10 $\mu$ M
Putidaredoxin reductase	1 $\mu$ M
Bovine liver catalase	20 $\mu$ g/ml
KCl	200 mM
Substrate	Typically 1 mM
NADH	250 - 400 $\mu$ M

\* 50 mM Tris-HCl buffer pH 7.4 is added to make up the volume.

\* Temperature controlled at 30°C, optional.

\* The NADH turnover rate could be determined by monitoring the absorbance at 340 nm with time.

\* Catalase does not catalyse the substrate oxidation reactions but rather it is present to remove any hydrogen peroxide by-product which could otherwise denature the P450<sub>cam</sub>.

The method can be increased in scale to, for example, 20 ml total incubation volume to allow purification of sufficient products by HPLC for

-28-

spectroscopic characterisation. Fresh substrate (1 mM) and NADH (1 - 2 mM) are added periodically, such as every 20 minutes in a total reaction time of, typically, 3 hours.

### Example 3

#### The in vivo system

The in vivo systems were expressed using the vector pGLW11, a derivative of the plasmid pKK223 (Brosius, J. and Holy, A. *Proc. Natl. Acad. Sci. USA*, 1984, 81, 6929-6933). Expression is directed by the *tac* promoter and the vector incorporates a gene conferring resistance to the antibiotic ampicillin.

Two systems were constructed. The first one expressed the electron transfer proteins putidaredoxin reductase (*camA* gene) and putidaredoxin (*camB* gene) as a fusion protein with a seven amino acid peptide linker, and the P450<sub>cam</sub> enzyme (*camC* gene) was expressed by the same vector but it was not fused to the electron proteins. The second system expressed the three proteins as separate entities in the *E. Coli* host. Both systems were catalytically competent for substrate oxidation in vivo.

The general strategy was as follows. The genes for the three proteins were cloned using *Eco* RI and *Hind* III as flanking sites, with *Eco* RI at the 5' end. For both in vivo systems there are restriction sites between the genes, including between the reductase and redoxin genes in the fusion construct. These restriction sites were introduced by PCR, as detailed below. The first task, however, was to carry out a silent mutation to remove the *Hind* III site within the *camA* gene for the reductase. The AAGCTT *Hind* III recognition sequence in the *camA* gene was changed to AAGCCT, which is a silent mutation because

-29-

GCT and GCC both encode alanine. The gene was completely sequenced to ensure that there were no spurious mutations.

## 1. The fusion protein system

### 1.a Manipulation of the *camA* gene by PCR

For the *camA* gene the primer below was used at the 5' end of the gene to introduce the *Eco* RI cloning site and to change the first codon from GTG to the strong start codon ATG.

5'- GAG ATT AAG AAT TCA TAA ACA CAT GGG AGT GCG TGC CAT ATG AAC GCA AAC

*Eco* RI

RBS

|→*camA*

At the 3' end of *camA* the primer was designed such that 15 bases are complementary to nucleotide sequence of the last five amino acid residues of *camA*. The stop codon immediately after the GCC codon for the last amino acid was removed, and then part of a seven amino acid linker (Thr Asp Gly Gly Ser Ser Ser) which contained a *Bam* HI cloning site (GGATCC = Gly Ser) was introduced. The coding sequence was thus:

5'- GAA CTG AGT AGT GCC ACT GAC GGA GGA TCC TCA TCG-3'

*camA* ~ Thr Asp Gly Gly Ser

|*Bam* HI|

The primer sequence shown below is the reverse complement used for PCR:

5'- CGA TGA GGA TCC TCC GTC AGT GGC ACT ACT CAG TTC-3'

### 1.b Manipulations of the *camB* gene by PCR

For the *camB* gene the primer at the 5' end incorporated the second half of the peptide linker

-30-

between the reductase and redoxin proteins, and the restriction site *Bam* HI for joining the two amplified genes together.

5'- TCA TCG **GGA TCC** TCA TCG **ATG** TCT AAA GTA GTG TAT-3'

Gly Ser Ser Ser |- *camB*

|*Bam* HI| Start

At the 3' end of *camB* the primer incorporates 12 nucleotides complementary to the end of *camB* followed by the stop codon TAA, a 6 nucleotide spacer before the GGAG ribosome binding site. *Xba* I and *Hind* III sites were then added to allow cloning of the *camC* gene when required. The sequence of the coding strand was therefore:

5'- CCC GAT AGG CAA TGG **TAA** TCA TCG **GGAG** TCT **AGA** GCA TCG **AAG** CTT TCA TCG-3'

*CamB* -|stop RBS *Xba* I *Hind* III

The primer shown below is the reverse complement used for PCR:

5'-CGA TGA AAG CTT CGA TGC TCT AGA CTCC CGA TGA TTA CCA TTG CCT ATC GGG -3'

### 1.c Preparation of the full fusion construct

The *camA* and *camB* genes were amplified by the PCR using the primers described above. The new *camA* was digested with *Eco* RI and *Bam* HI, while the new *CamB* was digested with *Bam* HI and *Hind* III. The pGLW11 expression vector was digested with *Eco* RI and *Hind* III. All three were purified by agarose gel electrophoresis and the three gel slices containing the separate fragments were excised from the gel and ligated together, and then transformed into *E. Coli* DH5 $\alpha$ . Successful ligation of all the fragments were confirmed by a series of restriction

-31-

digestion experiments, especially the presence of the new and unique *Xba* I site. The entire sequence of the insert from the *Eco* RI site to the *Hind* III site was determined to ensure that all the sequences were correct.

5 The new plasmid, named pSGB<sup>F</sup>, was transformed into *E. Coli* and expression of the reductase and redoxin proteins was induced by IPTG. When a purified P450<sub>cam</sub> enzyme was added to the cell-free extract, substrate oxidation was observed for a variety of substrates.

10 When the *camC* gene is cloned into the pSGB<sup>F</sup> plasmid using the *Xba* I and *Hind* III restriction sites, the new recombinant plasmid thus generated expresses the reductase and redoxin as a fusion protein and the P450<sub>cam</sub> enzyme as a separate entity both from the same mRNA molecule. This *in vivo* system is catalytically competent for terpene oxidation in whole cells.

20 2. The *in vivo* system with the protein expressed separately

2.a The basic strategy

25 The starting point of the preparation of this *in vivo* system was the recombinant plasmid used to express the *camA* gene for putidaredoxin reductase. The *camA* gene was cloned into the pGLW11 plasmid using the *Eco* RI and *Bam* HI restriction sites, with *Eco* RI being at the 5' end of the gene. Conveniently the polylinker region of the pGLW11 vector has a *Hind* III site downstream of the *Bam* HI site. The *camB* gene was therefore manipulated by PCR such that it can be cloned into pGLW11 using the *Bam* HI and *Hind* III sites. This new plasmid expresses the reductase and redoxin as separate proteins.

30 The *camB* gene was cloned into pUC118 by the *Bam* HI



-32-

and *Hind* III cloning sites to express putidaredoxin for our general *in vitro* substrate oxidation work.

Therefore, the PCR primer at the 3' end of the *camB* gene was designed to introduce a ribosome binding site and the *Xba* I restriction site upstream of the *Hind* III site so that the *camC* gene can be inserted downstream of *camB* using the *Xba* I and *Hind* III sites. Therefore the three genes were cloned without fusion in the pGLW11 expression vector and arranged in the order 5'-*camA-camB-camC*-3', and each gene has its own RBS to initiate protein synthesis.

## 2.b Manipulations of the *camB* gene

We used the internal and unique restriction site *Mlu* I (recognition sequence ACGCGT) within the *camB* gene as the starting point so that the PCR product has a different size from the PCR template fragment. The primers were as follows:

5'- TCA TCG **ACG CGT** CGC GAA CTG CTG-3'

where the *Mlu* I site is in bold.

The desired coding sequence at the 3' end of the *camB* gene was:

5'- CCC GAT AGG CAA TGG **TAA** GTA GGT GAA TAT CTA ATC CCC ATC  
*camB* -|stop

TAT GCG CGA GTG **GAG TCT** AGA GTT CGA-3'

RBS *Xba* I

After the stop codon there is a 35 base spacer before the RBS which is used to initiate the synthesis of the P450<sub>cam</sub> enzyme. The *Xba* I cloning site is located

-33-

within the spacer between the RBS and the start codon (not in this primer) of the *camC* gene. The PCR primer used was the reverse complement of the sequence above. The PCR was carried out and the amplified fragment of the appropriate size was purified by agarose gel electrophoresis and the gel slice excised.

One extra step was necessary to complete the construction of the new plasmid. The plasmid for the fusion protein *in vivo* system was digested with *Mlu* I and *Hind* III restriction enzymes, purified by agarose gel electrophoresis, and the gel slice for the small *camB* fragment excised. The pUC118 plasmid for *camB* expression was similarly digested, and the gel slice for the backbone was excised. By ligating the two fragments together we prepared a new pUC118-based plasmid which had an *Xba* I site followed by an *Hind* III site downstream of the stop codon of *camB*. This new plasmid was digested with the *Mlu* I and *Xba* I enzymes and the backbone was ligated with the new *camB* fragment described above to generate a plasmid with the following arrangement of the key components:

..lac Promoter..Bam HI..*camB* gene..spacer..RBS..Xba I..*Hind* III..

## 2.c Preparation of the *in vivo* system plasmid

Once the modified *camB* with the *Xba* I and *Hind* III restriction sites and appropriate spacers were prepared, the *in vivo* system was constructed by cloning this into the pGLW11-based plasmid used to express the *camA* gene (reductase protein) using the *Bam* HI and *Hind* III sites. The new *in vivo* system vector has the following arrangement of the key components:

..tac Promoter..Eco IRI..RBS..*camA* gene..spacer..Bam

-34-

HI..RBS..**camB gene**..spacer..RBS..*Xba* I.. *Hind* III..

This new plasmid, named pSGB<sup>+</sup>, was transformed into *E. Coli* and expression of the reductase and redoxin proteins was induced by IPTG. When a purified P450<sub>cam</sub> enzyme was added to the cell-free extract, substrate oxidation was observed for a variety of substrates.

When the *camC* gene is cloned into this pSGC<sup>+</sup> plasmid using the *Xba* I and *Hind* III restriction sites, the new recombinant plasmid thus generated will express the three proteins separately, each under the direction of its own RBS but from the same mRNA molecule. Thus constitutes the *in vivo* system used in the vast majority of our terpene oxidation work.

### 3. Introduction of an *Xba* I site into pRH1091

This is the final step to enable the *camC* gene to be cloned into the *in vivo* systems by the two cloning sites *Xba*I and *Hind* III. The *Xba* I site was added by PCR of the entire pRH1091 plasmid using two primers. The presence of these two sites will also enable cloning of the *camC* gene into M13 since both *Xba* I and *Hind* III are unique in *camC* and M13.

The primers shown below maintain the *Hind* III cloning site AAGCTT:

5'-TCA TCG **AAG CTT** GGC TGT TTT-3'

*Hind* III |→ vector

At the other end the coding sequence desired was:

5'-ACA ATT TCA CAC **AGGA TCT** AGA C **CAT ATG** TCA TCG **AAG CTT** TCA TCG-3'

Vector -|RBS *Xba* I      *Nde* I      *Hind* III

-35-

This sequence maintained the *Nde* I and *Hind* III sites but the new *Xba* I site was introduced upstream of the *Nde* I site. The PCR primer used was the reverse complement of the desired sequence:

5'-CGA TGA AAG CTT CGA TGA CAT ATG GTC T AGA TCCT GTG TGA AAT  
TGT-3'

The PCR product was then purified by agarose gel electrophoresis, digested with *Hind* III and circularised with T4 DNA ligase. Success of the PCR method was indicated by the presence of a new and unique *Xba* I site in plasmid DNA isolated from transformants.

#### 4. Cloning of *camC* into the *in vivo* systems

All existing *camC* mutants were cut out of pRH1091-based expression plasmids with *Nde* I and *Hind* III. The new vector is similarly cut with the same restriction enzymes and the *camC* gene cloned into this plasmid with T4 DNA ligase. This DNA is transformed into *E. Coli* JM109 which then may be grown to express P450<sub>cam</sub>.

The *camC* gene is excised from the new vector using *Xba* I and *Hind* III restriction enzymes and cloned into either the *in vivo* vector systems or M13mp19 for mutagenesis.

#### 5. *In vivo* expression and substrate turnover

For protein expression, cells are grown in LBamp medium (tryptone 10 g/litre, yeast extract 5 g/litre, NaCl 10 g/litre, 50 µg/ml ampicillin) at 30°C until the OD<sub>600nm</sub> reaches 1.0 - 1.2. IPTG (isopropyl-β-D-thiogalactopyranoside) was added to a final concentration of 1 µM (from a 1 M stock in H<sub>2</sub>O) and the culture was incubated at 30°C overnight.

-36-

For simple screening the substrate can be added to culture and the incubation continued. However, due to impurities from the culture media the cells were generally washed twice with 0.5 vol. of buffer P, ( $\text{KH}_2\text{PO}_4$  6.4 g,  $\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O}$  25.8 g,  $\text{H}_2\text{O}$  to 4 litres, pH 7.4) and resuspended in 0.25 vol. oxygen saturated buffer P containing 24 mM glucose. Substrate was added to 1 mM and the incubation continued at 30°C. The reaction was allowed to run for 24 hours with periodic additions of substrate and glucose.

The reaction was analysed by extracting 1 ml of the incubation mixture with 250  $\mu\text{l}$  of ethylacetate. After centrifuging in a microcentrifuge at 13,000 g for 2 minutes, 2  $\mu\text{l}$  of the organic extract was injected onto a 0.25 mm x 30 m DB-1 gas chromatography column in a Fisons GC 8000 series gas chromatograph. The samples were carried through the column using helium carrier gas and the compounds present were detected using a flame ionisation detector.

A variety of temperature programmes were used for different substrate to resolve turnover products.

#### Monoterpenes

Injector temperature	150 °C
Detector temperature	250 °C
Oven temperature	120 °C for 15 min → 200 °C at 25 °C/min, 200 °C for 1 min.

#### Sesquiterpenes

Injector temperature	250 °C
Detector temperature	250 °C
Oven temperature	150 °C → 230 °C at 5 °C/min, 230 °C for 1 min.

Results for particular oxidation reactions are shown

-37-

in table 6 and in the Figures.

Figure 1 shows the result of an oxidation reaction with camphor. The 5-ketocamphor arises from further oxidation of 5-exo-hydroxycamphor. As can be seen there is surprisingly little of the further oxidation occurring in the presence of camphor.

#### Example 4

##### A second in vivo expression system

The cluster of genes for the expression of the three proteins of the P450<sub>cam</sub> system, as described in Example 3, were also expressed in whole *E. coli* cells using the pCW vector. This vector utilises two *tac* promoters arranged in line to increase protein expression. It has a RBS, and contains the gene conferring resistance to the antibiotic ampicillin (Barnes, H.J. *Methods Enzymol.* 1996, 272, 3-14).

Both methods of expressing catalytically competent P450<sub>cam</sub> systems described in Example 3 were successful with the pCW vector. Thus the fusion system, where putidaredoxin reductase and putidaredoxin were expressed as a fusion protein with an oligopeptide linker, but the P450<sub>cam</sub> monooxygenase was expressed but not fused to the electron transfer proteins. The second system expressed all three proteins as separate entities in the same *E. coli* host.

##### **1. The fusion protein system**

A new plasmid was constructed by cloning the gene for the fusion of electron transfer proteins into pCW so that different P450<sub>cam</sub> mutants could be introduced into the system by cloning. The 5' end oligonucleotide used for the PCR amplification of the *cam A* gene introduced not only the *Eco* RI site for cloning into the pGL W11

-38-

vector but also a *Nde* I site which spans the ATG start codon of the gene (see Example 3, section 1a). In the pCW vector there is a *Nde* I site positioned downstream of the RBS for cloning of the gene to be expressed. The pGL W11 vector containing the *camA-camB* fusion gene was digested with *Nde* I and *Hind* III, and the insert purified by agarose gel electrophoresis. The pCW vector system was also digested with these two enzymes and the linearised vector purified by the same method. The two fragments were ligated with DNA ligase to generate the new pCWSGB<sub>F</sub> plasmid based on the pCW vector and which expressed the fusion of the electron transfer proteins. The insert excised from the pGL W11-based plasmid already contained a RBS for protein expression and an *Xba* I site just upstream of the *Hind* III site (see Example 3, section 2.c), so that the *cam C* gene encoding P450<sub>cam</sub> mutants can be cloned using these two sites and expressed off the RBS. Therefore genes encoding the P450<sub>cam</sub> mutants can be excised from the modified pRH system and cloned into the new pCWSGB<sub>F</sub> plasmid using the *Xba* I and *Hind* III sites.

## 2. The three proteins expressed separately

This system was generated in exactly the same way as for the fusion system. Thus the pSGB+ plasmid was digested with the *Nde* I and *Hind* III restriction enzymes and the insert cloned into the pCW vector. This new plasmid pCWSGB+ expressed putidaredoxin reductase and putidaredoxin as separate entities off the twin *tac* promoters of the pCW vector. The P450<sub>cam</sub> mutants were introduced into this vector using the *Xba* I and *Hind* III sites.

## 3. In vivo expression and substrate turnover

-39-

The conditions described below were used for test purposes in shake flasks in a laboratory and were not optimised. Under proper fermenter conditions with higher expression and biomass the final yield of products will be much increased.

*E. coli* DH5 $\alpha$  cells harbouring either of the catalytically competent P450<sub>cam</sub> systems were grown from a single colony on an agar plate in 1 L of LBamp medium at 30 °C until OD<sub>600nm</sub> reached ca. 1. IPTG was added to 1 mM final concentration (from a 1 M stock in water) and the culture grown for a further 6 h. The final OD<sub>600nm</sub> were in the range 2.0 - 2.5. Cells were harvested by centrifugation at 5000 g and washed once with 40 mM phosphate buffer, pH 7.4. The cell pellet was resuspended in 500 ml of 40 mM phosphate buffer, pH 7.4. Glucose was added as a 2 M stock to a final concentration of 100 mM, and 1 mL of the substrate ( $\alpha$ -pinene or R-limonene) was added to start the reaction. The mixture was shaken in an open 2 L conical flask in an orbital incubator at 200 rpm. More glucose was added every 24 h (100 mM final concentration based on a 500 mL volume, from a 2 M stock) and more substrate (1 mL) was added every 12 h. The progress of the reaction was monitored by GC and the whole cell system was active for at least 5 days at ambient temperatures and the minimum yield at the end of day 5, as assayed by extraction of the reaction medium with chloroform and analysing by GC, was 100 mg of products without accounting for volatilisation of products into the atmosphere and condensation on the flask above the liquid level. In addition, no compounds arising from further hydroxylation at another carbon atom were observed when the substrate was present.




-40-

**Example 5****In vitro and in vivo substrate oxidation by P450<sub>BM-3</sub>**


In a typical reaction *in vitro* (optional 30°C temperature control) the 1.5 mL incubation mixture contained 40 mM phosphate buffer, pH 8.0, 1  $\mu$ M P450<sub>BM-3</sub>, 50  $\mu$ g/mL of catalase, and terpene substrates were added as a 1 M stock in ethanol to a final concentration of 2 mM. NADPH was added typically to 400  $\mu$ M final concentration, and the rates of reaction could be monitored at 340 nm. After all the NADPH had been consumed, the mixture was extracted by vortexing with 0.5 mL of chloroform, the phases separated by centrifugation, and the organic phase could be analysed by GC using the programs described in Example 3.

Catalase does not catalyse the substrate oxidation reaction but rather it is present to remove any hydrogen by-product which could otherwise denature the enzyme. The method can be increased in scale to, for example, 20 mL total incubation volume to allow purification of sufficient produces by HPLC for spectroscopic characterisation. Fresh substrate (1 mM) and NADPH (1-2 mM) are added periodically, such as every 20 minutes in a total reaction time of, typically 3 hours.


Since the P450<sub>BM-3</sub> enzyme is catalytically self-sufficient, i.e., both the monooxygenase and electron transfer domains are in a single polypeptide, the enzyme as expressed in *E. coli* can be used for whole cell, *in vivo* substrate oxidations. The procedure described under Example 3 for *in vivo* substrate oxidation by the P450<sub>cam</sub> enzyme can also be used for the P450<sub>BM-3</sub> enzyme.




Limonene




α-pinene




β-pinene




α-terpinene




γ-terpinene




(+)-sabinene




(-)-α-thujene




myrcene



nerol



geraniol



ocimene

Table 1












Table 2

Table 3. HYDROPATHY SCALE FOR AMINO ACID  
SIDE CHAINS

Side Chain	Hydropathy
Ile	4.5
Val	4.2
Leu	3.8
Phe	2.8
Cys	2.5
Met	1.9
Ala	1.8
Gly	-0.4
Thr	-0.7
Ser	-0.8
Trp	-0.9
Tyr	-1.3
Pro	-1.6
His	-3.2
Glu	-3.5
Gln	-3.5
Asp	-3.5
Asn	-3.5
Lys	-3.9
Arg	-4.5

Table 4: P450<sub>cam</sub> mutants

All mutants optionally contain the base mutation C334A.

Single mutants: Y96A, Y96F, Y96L, Y96W.

Double mutants:

Y96A-F87A	Y96F-F87A	Y96F-V295A	Y96L-F87A	Y96L-A296L
Y96A-F87L	Y96F-F87I	Y96F-V295L	Y96L-F87L	Y96L-A296F
Y96A-F87W	Y96F-F87L	Y96F-V295I	Y96L-F98W	Y96L-V396A
Y96A-F98W	Y96F-F87W	Y96F-A296L	Y96L-T101L	Y96L-V396L
Y96A-L244A	Y96F-F98W	Y96F-A296F	Y96L-T101F	Y96L-V396F
Y96A-V247A	Y96F-T101L	Y96F-I395F	Y96L-L244A	Y96L-V396W
Y96A-V247L	Y96F-T101F	Y96F-I395G	Y96L-L244F	
Y96A-I395F	Y96F-T185A	Y96F-V396A	Y96L-V247A	
Y96A-I395G	Y96F-T185F	Y96F-V396L	Y96L-V247L	Y96W-F87W
	Y96F-T185L	Y96F-V396F	Y96L-V247F	Y96W-F98W
	Y96F-L244A	Y96F-V396W	Y96L-V247W	Y96W-L244A
	Y96F-V247A		Y96L-G248L	Y96W-V247A
	Y96F-V247L		Y96L-V295L	Y96W-V396A
	Y96F-G248L		Y96L-V295F	

Triple Mutants:

Y96A-F87A-L244A	Y96L-V247A-V396L	Y96F-F87W-V247A
Y96A-F87A-V247A	Y96L-V247A-V396F	Y96F-F87W-V247L
Y96A-F87L-L244A	Y96L-V247A-V396W	Y96F-F87W-V247F
Y96A-F87L-V247A	Y96L-V247F-V396A	Y96F-F87W-V295L
Y96A-L244A-V247A		Y96F-F87W-A296L
	Y96F-F87A-L244A	Y96F-F87W-V396A
Y96L-F87A-L244A	Y96F-F87A-V247A	Y96F-F87W-V396L
Y96L-F87A-V247A	Y96F-F87A-V247L	Y96F-V247F-V396A
Y96L-F87L-L244A	Y96F-F87A-I395F	Y96F-L244A-V396L
Y96L-F87L-V247A	Y96F-F87A-I395G	Y96F-L244A-V396F
Y96L-V247A-I395F	Y96F-F87L-V247A	Y96F-L244A-V396W
Y96L-V247L-I395F	Y96F-F87L-V247L	Y96F-L244F-V396A
Y96L-V247L-I395G	Y96F-F87L-I395F	Y96F-V247A-V396L
Y96L-L244A-V396L	Y96F-F87W-T185A	Y96F-V247A-V396F
Y96L-L244A-V396F	Y96F-F87W-T185F	Y96F-V247A-V396W
Y96L-L244A-V396W	Y96F-F87W-T185L	
Y96L-L244F-V396A	Y96F-F87W-L244F	Y96W-F87W-F98W

TOSTRO 6695860

Table 4 (continued)Four mutations:

Y96A-F87A-L244A-V247A

Y96A-F87L-L244A-V247A

Y96L-F87A-L244A-V247A

Y96L-F87L-L244A-V247A

Y96F-F87W-L244A-V295L

Y96F-F87W-L244F-V396A

Y96F-F87W-L244A-A296L

Y96F-F87W-V247A-V396L

Y96F-F87W-V247A-V396F

Y96F-F87W-V247L-V295A

Y96F-F87W-V247L-V396A

Y96F-F87W-V247F-V396A

Y96F-F87W-V247A-I395F

Y96F-F87W-V247L-I395G

Five mutations:

Y96F-F87W-T185L-V247L-V295L

Y96F-F87W-T185L-V247L-V396A

Y96F-F87W-T185L-V247L-V396L

Table 5: P450<sub>BM-3</sub> mutants

R47L-Y51F	R47L-Y51F-F42A	R47L-Y51F-F87A	R47L-Y51F-A264V	R47L-Y51F-I263F-A264L
R47A-Y51A	R47L-Y51F-F42L	R47L-Y51F-F87W	R47L-Y51F-A264L	R47L-Y51F-I263W-A264I
R47A-Y51L	R47A-Y51L-F42L	R47L-Y51F-I263A	R47L-Y51F-A264I	
		R47L-Y51F-I263F	R47L-Y51F-M354L	
		R47L-Y51F-I263L	R47L-Y51F-M354A	
		R47L-Y51F-I263W		
	R47L-Y51F-L75A-M354L		R47L-Y51F-L181A	
	R47A-Y51L-L75A-M354A		R47L-Y51F-L181W	
	R47L-Y51F-F87W-A264L			
	R47L-Y51F-F87W-A264I			
	R47L-Y51F-F87W-A264F			

Table 6: Summary of changes in selectivity of terpene oxidation by P450<sub>cam</sub> mutants. Approximate proportions of only the products arising from the insertion of a single oxygen atom, i.e., alcohols and epoxides, are given.

Mutant	Products, some also indicated by their retention times (min) on a 30 m DB-1 fused silica GC column						Comments
<b>L-Limonene</b>	5.23-5.33 min 1,2-Oxide	6.10-6.20 min Product	6.60 - 6.70 min Isopiperitenol	7.05-7.15 min Carveol	8.15-8.25 min Product		
Wild-type	-	30%	70%	-	-		Very small quantities
F87I-Y96F	8%	5%	5%	2%	80%		
Y96F-V247L	5%	-	>85%	5%	<5%		Other small peaks observed
Y96F-T185L	5%	<2%	>90%	<2%	<2%		Other small peaks observed
F87W-Y96F-V247L	<2%	-	>95%	<2%	-		
F87L-Y96F-V247L	5%	5%	5%	2%	>80%		
<b><math>\alpha</math>-Pinene</b>	Pinene oxide	Verbenol	Verbenone				
Wild-type	10%	30%	10%				Remaining 50% other products
Y96F-V247A	10%	50%	20%				Remaining 20% other products
Y96F	10%	30%	15%				Many other products
F87W-Y96F-V247L	-	70%	15%				Very few other products
<b>Valencene</b>	Nookatol region		Nookatone	Product	Product		
Wild-type	9.60-9.70 min	9.80-9.90 min	11.3-11.4 min	13.0-13.1 min	14.0-14.1 min		
Y96F-V247A	-	-	-	-	-		Many small peaks
Y96L-V247A	30%	15%	10%	<10%	>30%		
	15%	20%	10%	>40%	10%		
<b><math>\gamma</math>-Terpinene</b>	5.70-5.80 min	6.20-6.30 min					
Y96F	15%	85%					



Table 7: P450cam wild-type sequence

acg act gaa acc ata caa agc aac gcc aat ctt gcc cct ctg cca ccc 48  
 Thr Thr Glu Thr Ile Gln Ser Asn Ala Asn Leu Ala Pro Leu Pro Pro  
 1 5 10 15  
 cat gtg cca gag cac ctg gta ttc gac ttc gac atg tac aat ccg tcg 96  
 His Val Pro Glu His Leu Val Phe Asp Phe Asp Met Tyr Asn Pro Ser  
 20 25 30  
 aat ctg tct gcc ggc gtg cag gag gcc tgg gca gtt ctg caa gaa tca 144  
 Asn Leu Ser Ala Gly Val Gln Glu Ala Trp Ala Val Leu Gln Glu Ser  
 35 40 45  
 aac gta ccg gat ctg gtg tgg act cgc tgc aac ggc gga cac tgg atc 192  
 Asn Val Pro Asp Leu Val Trp Thr Arg Cys Asn Gly Gly His Trp Ile  
 50 55 60  
 gcc act cgc ggc caa ctg atc cgt gag gcc tat gaa gat tac cgc cac 240  
 Ala Thr Arg Gly Gln Leu Ile Arg Glu Ala Tyr Glu Asp Tyr Arg His  
 65 70 75 80  
 ttt tcc agc gag tgc ccg ttc atc cct cgt gaa gcc ggc gaa gcc tac 288  
 Phe Ser Ser Glu Cys Pro Phe Ile Pro Arg Glu Ala Gly Glu Ala Tyr  
 85 90 95  
 gac ttc att ccc acc tcg atg gat ccg ccc gag cag cgc cag ttt cgt 336  
 Asp Phe Ile Pro Thr Ser Met Asp Pro Pro Glu Gln Arg Gln Phe Arg  
 100 105 110  
 gcg ctg gcc aac caa gtg gtt ggc atg ccg gtg gtg gat aag ctg gag 384  
 Ala Leu Ala Asn Gln Val Val Gly Met Pro Val Val Asp Lys Leu Glu  
 115 120 125  
 aac cgg atc cag gag ctg gcc tgc tcg ctg atc gag agc ctg cgc ccg 432  
 Asn Arg Ile Gln Glu Leu Ala Cys Ser Leu Ile Glu Ser Leu Arg Pro  
 130 135 140  
 caa gga cag tgc aac ttc acc gag gac tac gcc gaa ccc ttc ccg ata 480  
 Gln Gly Gln Cys Asn Phe Thr Glu Asp Tyr Ala Glu Pro Phe Pro Ile  
 145 150 155 160

cgc atc ttc atg ctg ctc gca ggt cta ccg gaa gaa gat atc ccg cac 528  
 Arg Ile Phe Met Leu Leu Ala Gly Leu Pro Glu Glu Asp Ile Pro His  
 165 170 175  
 ttg aaa tac cta acg gat cag atg acc cgt ccg gat ggc agc atg acc 576  
 Leu Lys Tyr Leu Thr Asp Gln Met Thr Arg Pro Asp Gly Ser Met Thr  
 180 185 190  
 ttc gca gag gcc aag gag gcg ctc tac gac tat ctg ata ccg atc atc 624  
 Phe Ala Glu Ala Lys Glu Ala Leu Tyr Asp Tyr Leu Ile Pro Ile Ile  
 195 200 205  
 gag caa cgc agg cag aag ccg gga acc gac gct atc agc atc gtt gcc 672  
 Glu Gln Arg Arg Gln Lys Pro Gly Thr Asp Ala Ile Ser Ile Val Ala  
 210 215 220  
 aac ggc cag gtc aat ggg cga ccg atc acc agt gac gaa gcc aag agg 720  
 Asn Gly Gln Val Asn Gly Arg Pro Ile Thr Ser Asp Glu Ala Lys Arg  
 225 230 235 240  
 atg tgt ggc ctg tta ctg gtc ggc ggc ctg gat acg gtg gtc aat ttc 768  
 Met Cys Gly Leu Leu Leu Val Gly Gly Leu Asp Thr Val Val Asn Phe  
 245 250 255  
 ctc agc ttc agc atg gag ttc ctg gcc aaa agc ccg gag cat cgc cag 816  
 Leu Ser Phe Ser Met Glu Phe Leu Ala Lys Ser Pro Glu His Arg Gln  
 260 265 270  
 gag ctg atc gag cgt ccc gag cgt att cca gcc gct tgc gag gaa cta 864  
 Glu Leu Ile Glu Arg Pro Glu Arg Ile Pro Ala Ala Cys Glu Glu Leu  
 275 280 285  
 ctc cgg cgc ttc tcg ctg gtt gcc gat ggc cgc atc ctc acc tcc gat 912  
 Leu Arg Arg Phe Ser Leu Val Ala Asp Gly Arg Ile Leu Thr Ser Asp  
 290 295 300  
 tac gag ttt cat ggc gtg caa ctg aag aaa ggt gac cag atc ctg cta 960  
 Tyr Glu Phe His Gly Val Gln Leu Lys Lys Gly Asp Gln Ile Leu Leu  
 305 310 315 320

- 50 -

ccg cag atg ctg tct ggc ctg gat gag cgc gaa aac gcc tgc ccg atg 1008  
 Pro Gln Met Leu Ser Gly Leu Asp Glu Arg Glu Asn Ala Cys Pro Met  
 325 330 335  
 cac gtc gac ttc agt cgc caa aag gtt tca cac acc acc ttt ggc cac 1056  
 His Val Asp Phe Ser Arg Glu Lys Val Ser His Thr Thr Phe Gly His  
 340 345 350  
 ggc agc cat ctg tgc ctt ggc cag cac ctg gcc cgc cgg gaa atc atc 1104  
 Gly Ser His Leu Cys Leu Gly Gln His Leu Ala Arg Arg Glu Ile Ile  
 355 360 365  
 gtc acc ctc aag gaa tgg ctg acc agg att cct gac ttc tcc att gcc 1152  
 Val Thr Leu Lys Glu Trp Leu Thr Arg Ile Pro Asp Phe Ser Ile Ala  
 370 375 380  
 ccg ggt gcc cag att cag cac aag agc ggc atc gtc agc ggc gtg cag 1200  
 Pro Gly Ala Gln Ile Gln His Lys Ser Gly Ile Val Ser Gly Val Gln  
 385 390 395 400  
 gca ctc cct ctg gtc tgg gat ccg gcg act acc aaa gcg gta 1242  
 Ala Leu Pro Leu Val Trp Asp Pro Ala Thr Thr Lys Ala Val  
 405 410 414

Table 8: P450<sub>BK-3</sub> sequence

1 / 1 31 / 11  
 atg aca att aaa gaa atg cct cag cca aaa acg ttt gga gag ctt aaa aat tta ccg tta  
 Met Thr Ile Lys Glu Met Pro Gln Pro Lys Thr Phe Gly Glu Leu Lys Asn Leu Pro Leu  
 61 / 21 91 / 31  
 tta aac aca gat aaa ccg gtt caa gct ttg atg aaa att gcg gat gaa tta gga gaa atc  
 Leu Asn Thr Asp Lys Pro Val Gln Ala Leu Met Lys Ile Ala Asp Glu Leu Gly Glu Ile  
 121 / 41 151 / 51  
 ttt aaa ttc gag gcg cct ggt cgt gta acg cgc tac tta tca agt cag cgt cta att aaa  
 Phe Lys Phe Glu Ala Pro Gly Arg Val Thr Arg Tyr Leu Ser Ser Gln Arg Leu Ile Lys  
 181 / 61 211 / 71  
 gaa gca tgc gat gaa tca cgc ttt gat aaa aac tta agt caa gcg ctt aaa ttt gta cgt  
 Glu Ala Cys Asp Glu Ser Arg Phe Asp Lys Asn Leu Ser Gln Ala Leu Lys Phe Val Arg  
 241 / 81 271 / 91  
 gat ttt gca gga gac ggg tta ttt aca agc tgg acg cat gaa aaa aat tgg aaa aaa gcg  
 Asp Phe Ala Gly Asp Gly Leu Phe Thr Ser Trp Thr His Glu Lys Asn Trp Lys Lys Ala  
 301 / 101 331 / 111  
 cat aat atc tta ctt cca agc ttc agt cag cga gca atg aaa ggc tat cat gcg atg atg  
 His Asn Ile Leu Leu Pro Ser Phe Ser Gln Gln Ala Met Lys Gly Tyr His Ala Met Met  
 361 / 121 391 / 131  
 gtc gat atc gcc gtg cag ctt gtt caa aag tgg gag cgt cta aat gca gat gag cat att  
 Val Asp Ile Ala Val Gln Leu Val Gln Lys Trp Glu Arg Leu Asn Ala Asp Glu His Ile  
 421 / 141 451 / 151  
 gaa gta ccg gaa gac atg aca cgt tta acg ctt gat aca att ggt ctt tgc ggc ttt aac  
 Glu Val Pro Glu Asp Met Thr Arg Leu Thr Leu Asp Thr Ile Gly Leu Cys Gly Phe Asn  
 481 / 161 511 / 171  
 tat cgc ttt aac agc ttt tac cga gat cag cct cat cca ttt att aca agt atg gtc cgt  
 Tyr Arg Phe Asn Ser Phe Tyr Arg Asp Gln Pro His Pro Phe Ile Thr Ser Met Val Arg  
 541 / 181 571 / 191  
 gca ctg gat gaa gca atg aac aag ctg cag cga gca aat cca gac gac cca gct tat gat  
 Ala Leu Asp Glu Ala Met Asn Lys Leu Gln Arg Ala Asn Pro Asp Asp Pro Ala Tyr Asp  
 601 / 201 631 / 211  
 gaa aac aag cgc cag ttt caa gaa gat atc aag gtg atg aac gac cta gta gat aaa att  
 Glu Asn Lys Arg Gln Phe Gln Glu Asp Ile Lys Val Met Asn Asp Leu Val Asp Lys Ile  
 661 / 221 691 / 231  
 att gca gat cgc aaa gca agc ggt gaa caa agc gat gat tta tta acg cat atg cta aac  
 Ile Ala Asp Arg Lys Ala Ser Gly Glu Gln Ser Asp Asp Leu leu Thr His Met Leu Asn  
 721 / 241 751 / 251  
 gga aaa gat cca gaa acg ggt gag ccg ctt gat gac gag aac att cgc tat caa att att  
 Gly Lys Asp Pro Glu Thr Gly Glu Pro Leu Asp Asp Glu Asn Ile Arg Tyr Gln Ile Ile  
 781 / 261 811 / 271  
 aca ttc tta att gcg gga cac gaa aca aca agt ggt ctt tta tca ttt gcg ctg tat ttc  
 Thr Phe Leu Ile Ala Gly His Glu Thr Thr Ser Gly Leu Leu Ser Phe Ala leu Tyr Phe  
 841 / 281 871 / 291  
 tta gtg aaa aat cca cat gta tta caa aaa gca gca gaa gaa gca gca cga gtt cta gta  
 Leu Val Lys Asn Pro His Val Leu Gln Lys Ala Ala Glu Glu Ala Ala Arg Val Leu Val  
 901 / 301 931 / 311  
 gat cct gct cca agc tac aaa caa gtc aaa cag ctt aaa tat gtc ggc atg gtc tta aac  
 Asp Pro Val Pro Ser Tyr Lys Gln Val Lys Gln Leu Lys Tyr Val Gly Met Val Leu Asn  
 961 / 321 991 / 331  
 gaa gcg ctg cgc tta tgg cca act gct cct gcg ttt tcc cta tat gca aaa gaa gat acg  
 Glu Ala Leu Arg Leu Trp Pro Thr Ala Pro Ala Phe Ser Leu Tyr Ala Lys Glu Asp Thr  
 1021 / 341 1051 / 351  
 gtg ctt gga gga gaa tat cct tta gaa aaa ggc gac gaa cta atg gtt ctg att cct cag  
 Val Leu Gly Gly Glu Tyr Pro Leu Glu Lys Gly Asp Glu Leu Met Val Leu Ile Pro Gln  
 1081 / 361 1111 / 371  
 ctt cac cgt gat aaa aca att tgg gga gac gat gtg gaa gag ttc cgt cca gag cgt ttt  
 Leu His Arg Asp Lys Thr Ile Trp Gly Asp Asp Val Glu Glu Phe Arg Pro Glu Arg Phe  
 1141 / 381 1171 / 391  
 gaa aat cca agt gcg att ccg cag cat gcg ttt aaa ccg ttt gga aac ggt cag cgt gcg  
 Glu Asn Pro Ser Ala Ile Pro Gln His Ala Phe Lys Pro Phe Gly Asn Gly Gln Arg Ala  
 1201 / 401 1231 / 411  
 tgt atc ggt cag cag ttc gct ctt cat gaa gca acg ctg gta ctt ggt atg atg cta aaa  
 Cys Ile Gly Gln Gln Phe Ala Leu His Glu Ala Thr Leu Val Leu Gly Met Met Leu Lys  
 1261 / 421 1291 / 431  
 cac ttt gac ttt gaa gat cat aca aac tac gag ctg gat att aaa gaa act tta acg tta  
 His Phe Asp Phe Glu Asp His Thr Asn Tyr Glu Leu Asp Ile Lys Glu Thr Leu Thr Leu  
 1321 / 441 1351 / 451  
 aaa cct gaa ggc ttt gtg gta aaa gca aaa tcg aaa aaa att ccg ctt ggc ggt att cct  
 Lys Pro Glu Gly Phe Val Val Lys Ala Lys Ser Lys Lys Ile Pro Leu Gly Gly Ile Pro  
 1381 / 461 1411 / 471

- 53 -

```

cat gaa gac tat ctg tat caa gaa gag ctt gaa aac gcc caa agc gaa ggc atc att acg
His Glu Asp Tyr Leu Tyr Gln Glu Glu Leu Glu Asn Ala Gln Ser Glu Gly Ile Ile Thr
2881 / 961 2911 / 971
ctt cat acc gct ttt tct cgc atg cca aat cag ccg aaa aca tac gtt cag cac gta atg
Leu His Thr Ala Phe Ser Arg Met Pro Asn Gln Pro Lys Thr Tyr Val Gln His Val Met
2941 / 981 2971 / 991
gaa caa gac ggc aag aaa ttg att gaa ctt ctt gat caa gga gcg cac ttc tat att tgc
Glu Gln Asp Gly Lys Lys Leu Ile Glu Leu Leu Asp Gln Gly Ala His Phe Tyr Ile Cys
3001 / 1001 3031 / 1011
gga gac gga agc caa atg gca cct gcc gtt gaa gca acg ctt atg aaa agc tat gct gac
Gly Asp Gly Ser Gln Met Ala Pro Ala Val Glu Ala Thr Leu Met Lys Ser Tyr Ala Asp
3061 / 1021 3091 / 1031
gtt cac caa gtg agt gaa gca gac gct cgc tta tgg ctg cag cag cta gaa gaa aaa ggc
Val His Gln Val Ser Glu Ala Asp Ala Arg Leu Trp Leu Gln Gln Leu Glu Glu Lys Gly
3121 / 1041
cga tac gca aaa gac gtg tgg gct ggg taa
Arg Tyr Ala Lys Asp Val Trp Ala Gly OCH

```

105180\*66695660

tca cct agc act gaa cag tct gcc aaa aaa gca cgc aaa aag gca gaa aac gct cat aat  
 Ser Pro Ser Thr Glu Gln Ser Ala Lys Lys Val Arg Lys Lys Ala Glu Asn Ala His Asn  
 1441 / 481 1471 / 491  
 acg cgc ctg ctt gtg cta tac ggt tca aat atg gga aca gct gaa gga acg gcg cgt gat  
 Thr Pro Leu Leu Val Leu Tyr Gly Ser Asn Met Gly Thr Ala Glu Gly Thr Ala Arg Asp  
 1501 / 501 1531 / 511  
 tta gca gat att gca atg agc aaa gga ttt gca cgc cag gtc gca acg ctt gat tca cac  
 Leu Ala Asp Ile Ala Met Ser Lys Gly Phe Ala Pro Gln Val Ala Thr Leu Asp Ser His  
 1561 / 521 1591 / 531  
 gcc gga aat ctt ccg cgc gaa gga gct gta tta att gta acg gcg tct tat aac ggt cat  
 Ala Gly Asn Leu Pro Arg Glu Gly Ala Val Leu Ile Val Thr Ala Ser Tyr Asn Gly His  
 1621 / 541 1651 / 551  
 ccg cct gat aac gca aag caa ttt gtc gac tgg tta gac caa gcg tct gct gat gaa gta  
 Pro Pro Asp Asn Ala Lys Gln Phe Val Asp Trp Leu Asp Gln Ala Ser Ala Asp Glu Val  
 1681 / 561 1711 / 571  
 aaa ggc gtt cgc tac tcc gta ttt gga tgc ggc gat aaa aac tgg gct act acg tat caa  
 Lys Gly Val Arg Tyr Ser Val Phe Gly Cys Gly Asp Lys Asn Trp Ala Thr Thr Tyr Gln  
 1741 / 581 1771 / 591  
 aaa gtg cct gct ttt atc gat gaa acg ctt gcc gct aaa ggg gca gaa aac atc gct gac  
 Lys Val Pro Ala Phe Ile Asp Glu Thr Leu Ala Ala Lys Gly Ala Glu Asn Ile Ala Asp  
 1801 / 601 1831 / 611  
 cgc ggt gaa gca gat gca agc gac gac ttt gaa ggc aca tat gaa gaa tgg cgt gaa cat  
 Arg Gly Glu Ala Asp Ala Ser Asp Asp Phe Glu Gly Thr Tyr Glu Glu Trp Arg Glu His  
 1861 / 621 1891 / 631  
 atg tgg agt gac gta gca gcc tac ttt aac ctc gac att gaa aac agt gaa gat aat aaa  
 Met Trp Ser Asp Val Ala Ala Tyr Phe Asn Leu Asp Ile Glu Asn Ser Glu Asp Asn Lys  
 1921 / 641 1951 / 651  
 tct act ctt tca ctt caa ttt gtc gac agc gcc gcg gat atg ccg ctt gcg aaa atg cac  
 Ser Thr Leu Ser Leu Gln Phe Val Asp Ser Ala Ala Asp Met Pro Leu Ala Lys Met His  
 1981 / 661 2011 / 671  
 ggt cgc ttt tca acg aac gtc gta gca agc aaa gaa ctt caa cag cca ggc agt gca cga  
 Gly Ala Phe Ser Thr Asn Val Val Ala Ser Lys Glu Leu Gln Gln Pro Gly Ser Ala Arg  
 2041 / 681 2071 / 691  
 agc acg cga cat ctt gaa att gaa ctt cca aaa gaa gct tct tat caa gaa gga gat cat  
 Ser Thr Arg His Leu Glu Ile Glu Leu Pro Lys Glu Ala Ser Tyr Gln Glu Gly Asp His  
 2101 / 701 2131 / 711  
 tta ggt gtt att cct cgc aac tat gaa gga ata gta aac cgt gta aca gca agg ttc ggc  
 Leu Gly Val Ile Pro Arg Asn Tyr Glu Gly Ile Val Asn Arg Val Thr Ala Arg Phe Gly  
 2161 / 721 2191 / 731  
 cta gat gca tca cag caa atc cgt ctg gaa gca gaa gaa gaa aaa tta gct cat ttg cca  
 Leu Asp Ala Ser Gln Gln Ile Arg Leu Glu Ala Glu Glu Glu Lys Leu Ala His Leu Pro  
 2221 / 741 2251 / 751  
 ctc gct aaa aca gta tcc gta gaa gag ctt ctg caa tac gtg gag ctt caa gat cct gtt  
 Leu Ala Lys Thr Val Ser Val Glu Glu Leu Leu Gln Tyr Val Glu Leu Gln Asp Pro Val  
 2281 / 761 2311 / 771  
 acg cgc acg cag ctt cgc gca atg gct gct aaa acg gtc tgc ccg ccg cat aaa gta gag  
 Thr Arg Thr Gln Leu Arg Ala Met Ala Ala Lys Thr Val Cys Pro Pro His Lys Val Glu  
 2341 / 781 2371 / 791  
 ctt gaa gcc ttg ctt gaa aag caa gcc tac aaa gaa caa gtg ctg gca aaa cgt tta aca  
 Leu Glu Ala Leu Leu Glu Lys Gln Ala Tyr Lys Glu Gln Val Leu Ala Lys Arg Leu Thr  
 2401 / 801 2431 / 811  
 atg ctt gaa ctg ctt gaa aaa tac ccg gcg tgt gaa atg aaa ttc agc gaa ttt atc gcc  
 Met Leu Glu Leu Leu Glu Lys Tyr Pro Ala Cys Glu Met Lys Phe Ser Glu Phe Ile Ala  
 2461 / 821 2491 / 831  
 ctt ctg cca agc ata cgc ccg cgc tat tac tgc att tct tca tca cct cgt gtc gat gaa  
 Leu Leu Pro Ser Ile Arg Pro Arg Tyr Tyr Ser Ile Ser Ser Ser Pro Arg Val Asp Glu  
 2521 / 841 2551 / 851  
 aaa caa gca agc atc acg gtc agc gtt gtc tca gga gaa gcg tgg agc gga tat gga gaa  
 Lys Gln Ala Ser Ile Thr Val Ser Val Val Ser Gly Glu Ala Trp Ser Gly Tyr Gly Glu  
 2581 / 861 2611 / 871  
 tat aaa gga att gcg tgc aac tat ctt gcc gag ctg caa gaa gga gat acg att acg tgc  
 Tyr Lys Gly Ile Ala Ser Asn Tyr Leu Ala Glu Leu Gln Glu Gly Asp Thr Ile Thr Cys  
 2641 / 881 2671 / 891  
 ttt att tcc aca ccg cag tca gaa ttt acg ctg cca aaa gac cct gaa acg ccg ctt atc  
 Phe Ile Ser Thr Pro Gln Ser Glu Phe Thr Leu Pro Lys Asp Pro Glu Thr Pro Leu Ile  
 2701 / 901 2731 / 911  
 atg gtc gga ccg gga aca ggc gtc gcg ccg ttt aga ggc ttt gtg cag gcg cgc aaa cag  
 Met Val Gly Pro Gly Thr Gly Val Ala Pro Phe Arg Gly Phe Val Gln Ala Arg Lys Gln  
 2761 / 921 2791 / 931  
 cta aaa gaa caa gga cag tca ctt gga gaa gca cat tta tac ttc ggc tgc cgt tca cct  
 Leu Lys Glu Gln Gly Gln Ser Leu Gly Glu Ala His Leu Tyr Phe Gly Cys Arg Ser Pro  
 2821 / 941 2851 / 951

TIME: 10:15:47

Output Set: N:\CRF3\11212001\I856339A.raw

# ENTERED

11/27/01

## RAW SEQUENCE LISTING

DATE: 11/27/2001

PATENT APPLICATION: US/09/856,339A

TIME: 10:15:47

Input Set : A:\Sequence Listing.txt

Output Set: N:\CRF3\11212001\I856339A.raw

```

84 <220> FEATURE:
85 <221> NAME/KEY: misc_feature
86 <222> LOCATION: (1)..(3150)
87 <223> OTHER INFORMATION: Coding sequence
90 <400> SEQUENCE: 2
91 atgacaatta aagaaatgcc tcagccaaaa acgtttggag agcttaaaaa tttaccgtta      60
93 ttaaacacag ataaaccggt tcaagctttg atgaaaattg cggatgaatt aggagaaatc      120
95 tttaaattcg aggcgcctgg tcgtgtaacg cgctacttat caagtcagcg totaattaaa      180
97 gaagcatgcg atgaatcacg ctttgataaa aacttaagtc aagcgcttaa atttgtacgt      240
99 gattttgcag gagacgggtt atttacaagc tggacgcagc aaaaaaattg gaaaaaagcg      300
101 cataatatct tacttccaag cttcagtcag caggcaatga aaggctatca tgcgatgatg      360
103 gtcgatatcg ccgtgcagct tgttcaaaaag tgggagcgtc taaatgcaga tgagcatatt      420
105 gaagtaccgg aagacatgac acgtttaacg cttgatacaa ttggtctttg cggctttaac      480
107 tatcgcttta acagctttta ccgagatcag cctcatccat ttattacaag tatggtccgt      540
109 gcaactggatg aagcaatgaa caagctgcag cgagcaaatc cagacgaccc agcttatgat      600
111 gaaaacaagc gccagtttca agaagatatc aaggtgatga acgacctagt agataaaatt      660
113 attgcagatc gcaaagcaag cggtgaacaa agcgatgatt tattaacgca tatgctaaac      720
115 ggaaaagatc cagaaacggg tgagccgctt gatgacgaga acattcgcta tcaaattatt      780
117 acattcttaa ttgcgggaca cgaacaacaa agtggctctt tatcatttgc gctgtatttc      840
119 ttagtgaaaa atccacatgt attacaaaaa gcagcagaag aagcagcacg agttctagta      900
121 gatcctgctc caagctacaa acaagtcaaa cagcttaaat atgtcggcat ggtcttaaac      960
123 gaagcgctgc gcttatggcc aactgctcct gcgttttccc tatatgcaaa agaagatacg      1020
125 gtgcttgagg gagaatatcc tttagaaaaa ggcgacgaac taatggttct gattcctcag      1080
127 cttaaccgtg ataaaacaat ttggggagac gatgtggaag agttccgtcc agagcgtttt      1140
129 gaaaatccaa gtgcgattcc gcagcatgcg tttaaaccgt ttggaaaacg tcagcgctgcg      1200
131 tgtatcggtc agcagttcgc tcttcatgaa gcaacgctgg tacttggtat gatgctaaaa      1260
133 cactttgact ttgaagatca tacaaactac gagctggata ttaaagaaac tttaacgtta      1320
135 aaacctgaag gctttgtggt aaaagcaaaa tcgaaaaaaa ttccgcttgg cggatttcct      1380
137 tcacctagca ctgaacagtc tgccaaaaaa gcacgcaaaa aggcagaaaa cgctcataat      1440
139 acgccgctgc ttgtgctata cgtttcaaat atgggaacag ctgaaggaac ggcgcgtgat      1500
141 ttagcagata ttgcaatgag caaaggattt gcaccgcagg tcgcaacgct tgattcacac      1560
143 gccggaaatc ttccgcgcga aggagctgta ttaattgtaa cggcgtctta taacggtcat      1620
145 ccgcctgata acgcaaagca atttgtcgac tggttagacc aagcgtctgc tgatgaagta      1680
147 aaaggcgttc gtaactccgt atttggatgc ggcgataaaa actgggctac tacgtatcaa      1740
149 aaagtgcctg cttttatcga tgaaacgctt gccgctaaag ggcagaaaa catcgctgac      1800
151 cgcggtgaag cagatgcaag cgacgacttt gaaggcacat atgaagaatg gcgtgaacat      1860
153 atgtggagtg acgtagcagc ctactttaac ctgcacattg aaaacagtga agataataaa      1920
155 tctactcttt cacttcaatt tgtcgacagc gccgcggata tgccgcttgc gaaaatgcac      1980
157 ggtgcgtttt caacgaacgt cgtagcaagc aaagaacttc aacagccagg cagtgcacga      2040
159 agcacgcgac atcttgaaat tgaacttcca aaagaagctt cttatcaaga aggagatcat      2100
161 ttaggtgtta ttctcgcaa ctatgaagga atagtaaacc gtgtaacagc aaggttcggc      2160
163 ctatagtcac cacagcaaat ccgtctggaa gcagaagaag aaaaattagc tcatttgcca      2220
165 ctgcctaaaa cagtatccgt agaagagctt ctgcaatacg tggagcttca agatcctgtt      2280
167 acgcgcacgc agcttcgcgc aatggctgct aaaacggtct gccgcgcgca taaagtagag      2340
169 cttgaagcct tgcttgaaaa gcaagcctac aaagaacaag tgctggcaaa acgtttaaca      2400
171 atgcttgaac tgcttgaaaa ataocggcg tgtgaaatga aattcagcga atttatcgcc      2460
173 cttctgccaa gcatacggcc gcgtatttac tcgatttctt catcacctcg tgtcgatgaa      2520
175 aaacaagcaa gcatcacggt cagcgttgct tcaggagaag cgtggagcgg atatggagaa      2580
177 tataaaggaa ttgcgtcgaa ctatcttgcc gagctgcaag aaggagatac gattacgtgc      2640

```



## RAW SEQUENCE LISTING

DATE: 11/27/2001

PATENT APPLICATION: US/09/856,339A

TIME: 10:15:48

Input Set : A:\Sequence Listing.txt

Output Set: N:\CRF3\11212001\I856339A.raw

```

179 tttattttcca caccgcagtc agaatttacg ctgccaaaag accctgaaac gccgcttatc 2700
181 atggtcggac cggaacagg cgtcgcccg tttagaggct ttgtgcaggc gcgcaaacag 2760
183 ctaaaagaac aaggacagtc acttgagaa gcacatttat acttcggctg ccgttcacct 2820
185 catgaagact atctgtatca agaagagctt gaaaacgccc aaagcgaagg catcattacg 2880
187 cttcataccg cttttttctcg catgccaaat cagccgaaaa catacgttca gcacgtaatg 2940
189 gaacaagacg gcaagaaatt gattgaactt cttgatcaag gagcgcaact ctatatattgc 3000
191 ggagacggaa gccaaatggc acctgccgtt gaagcaacgc ttatgaaaag ctatgctgac 3060
193 gttcaccaag tgagtgaagc agacgctcgc ttatggctgc agcagctaga agaaaaaggc 3120
195 cgatacgcaa aagacgtgtg ggctgggtaa 3150

```

198 &lt;210&gt; SEQ ID NO: 3

199 &lt;211&gt; LENGTH: 7

200 &lt;212&gt; TYPE: PRT

201 &lt;213&gt; ORGANISM: Artificial Sequence

203 &lt;220&gt; FEATURE:

204 &lt;221&gt; NAME/KEY: DOMAIN

205 &lt;222&gt; LOCATION: (1)..(7)

206 &lt;223&gt; OTHER INFORMATION: Synthetic linker

209 &lt;400&gt; SEQUENCE: 3

211 Thr Asp Gly Thr Ser Ser Thr

212 1 5

214 &lt;210&gt; SEQ ID NO: 4

215 &lt;211&gt; LENGTH: 7

216 &lt;212&gt; TYPE: PRT

217 &lt;213&gt; ORGANISM: Artificial Sequence

219 &lt;220&gt; FEATURE:

220 &lt;221&gt; NAME/KEY: DOMAIN

221 &lt;222&gt; LOCATION: (1)..(7)

222 &lt;223&gt; OTHER INFORMATION: Synthetic Linker

225 &lt;400&gt; SEQUENCE: 4

227 Thr Asp Gly Ala Ser Ser Ser

228 1 5

230 &lt;210&gt; SEQ ID NO: 5

231 &lt;211&gt; LENGTH: 17

232 &lt;212&gt; TYPE: PRT

233 &lt;213&gt; ORGANISM: Artificial Sequence

235 &lt;220&gt; FEATURE:

236 &lt;221&gt; NAME/KEY: DOMAIN

237 &lt;222&gt; LOCATION: (1)..(17)

238 &lt;223&gt; OTHER INFORMATION: Synthetic Linker

241 &lt;400&gt; SEQUENCE: 5

243 Thr Asp Gly Thr Arg Pro Gly Pro Gly Pro Gly Pro Gly Pro Ser Ser

244 1 5 10 15

246 Thr

249 &lt;210&gt; SEQ ID NO: 6

250 &lt;211&gt; LENGTH: 21

251 &lt;212&gt; TYPE: PRT

252 &lt;213&gt; ORGANISM: Artificial Sequence

254 &lt;220&gt; FEATURE:

255 &lt;221&gt; NAME/KEY: DOMAIN

## RAW SEQUENCE LISTING

DATE: 11/27/2001

PATENT APPLICATION: US/09/856,339A

TIME: 10:15:48

Input Set : A:\Sequence Listing.txt

Output Set: N:\CRF3\11212001\I856339A.raw

256 <222> LOCATION: (1)..(21)  
 257 <223> OTHER INFORMATION: Synthetic Linker  
 260 <400> SEQUENCE: 6  
 262 Thr Asp Gly Thr Arg Pro Gly Pro Gly Pro Gly Pro Gly Pro Gly Pro  
 263 1 5 10 15  
 265 Gly Pro Ser Ser Thr  
 266 20  
 268 <210> SEQ ID NO: 7  
 269 <211> LENGTH: 4  
 270 <212> TYPE: PRT  
 271 <213> ORGANISM: Artificial Sequence  
 273 <220> FEATURE:  
 274 <221> NAME/KEY: DOMAIN  
 275 <222> LOCATION: (1)..(4)  
 276 <223> OTHER INFORMATION: Synthetic Linker  
 279 <400> SEQUENCE: 7  
 281 Pro Leu Glu Leu  
 282 1  
 284 <210> SEQ ID NO: 8  
 285 <211> LENGTH: 7  
 286 <212> TYPE: PRT  
 287 <213> ORGANISM: Artificial Sequence  
 289 <220> FEATURE:  
 290 <221> NAME/KEY: DOMAIN  
 291 <222> LOCATION: (1)..(7)  
 292 <223> OTHER INFORMATION: Synthetic Linker  
 295 <400> SEQUENCE: 8  
 297 Thr Asp Gly Gly Ser Ser Ser  
 298 1 5  
 300 <210> SEQ ID NO: 9  
 301 <211> LENGTH: 51  
 302 <212> TYPE: DNA  
 303 <213> ORGANISM: Artificial Sequence  
 305 <220> FEATURE:  
 306 <221> NAME/KEY: misc\_feature  
 307 <222> LOCATION: (1)..(51)  
 308 <223> OTHER INFORMATION: Synthetic Primer  
 311 <400> SEQUENCE: 9  
 312 gagattaaga attcataaac acatgggagt gcgtgccata tgaacgcaaa c 51  
 315 <210> SEQ ID NO: 10  
 316 <211> LENGTH: 36  
 317 <212> TYPE: DNA  
 318 <213> ORGANISM: Artificial Sequence  
 320 <220> FEATURE:  
 321 <221> NAME/KEY: variation  
 322 <222> LOCATION: (1)..(36)  
 323 <223> OTHER INFORMATION: Synthetic desired coding sequence  
 326 <400> SEQUENCE: 10  
 327 gaactgagta gtgccactga cggaggatcc tcatcg 36

## RAW SEQUENCE LISTING

DATE: 11/27/2001

PATENT APPLICATION: US/09/856,339A

TIME: 10:15:48

Input Set : A:\Sequence Listing.txt

Output Set: N:\CRF3\11212001\I856339A.raw

```

330 <210> SEQ ID NO: 11
331 <211> LENGTH: 36
332 <212> TYPE: DNA
333 <213> ORGANISM: Artificial Sequence
335 <220> FEATURE:
336 <221> NAME/KEY: misc_feature
337 <222> LOCATION: (1)..(36)
338 <223> OTHER INFORMATION: Synthetic Primer
341 <400> SEQUENCE: 11
342 cgatgaggat cctccgtcag tggcactact cagttc          36
345 <210> SEQ ID NO: 12
346 <211> LENGTH: 36
347 <212> TYPE: DNA
348 <213> ORGANISM: ARTIFICIAL SEQUENCE
350 <220> FEATURE:
351 <221> NAME/KEY: misc_feature
352 <222> LOCATION: (1)..(36)
353 <223> OTHER INFORMATION: Synthetic Primer
356 <400> SEQUENCE: 12
357 tcatcgggat cctcatcgat gtctaaagta gtgtat          36
360 <210> SEQ ID NO: 13
361 <211> LENGTH: 52
362 <212> TYPE: DNA
363 <213> ORGANISM: Artificial Sequence
365 <220> FEATURE:
366 <221> NAME/KEY: misc_feature
367 <222> LOCATION: (1)..(52)
368 <223> OTHER INFORMATION: Synthetic desired coding sequence
371 <400> SEQUENCE: 13
372 cccgataggc aatggtaatc atcgggagtc tagagcatcg aagctttcat cg          52
375 <210> SEQ ID NO: 14
376 <211> LENGTH: 52
377 <212> TYPE: DNA
378 <213> ORGANISM: Artificial Sequence
380 <220> FEATURE:
381 <221> NAME/KEY: misc_feature
382 <222> LOCATION: (1)..(52)
383 <223> OTHER INFORMATION: Synthetic Primer
386 <400> SEQUENCE: 14
387 cgatgaaagc ttcatgtctc tagactcccg atgattacca ttgcctatcg gg          52
390 <210> SEQ ID NO: 15
391 <211> LENGTH: 24
392 <212> TYPE: DNA
393 <213> ORGANISM: Artificial Sequence
395 <220> FEATURE:
396 <221> NAME/KEY: misc_feature
397 <222> LOCATION: (1)..(24)
398 <223> OTHER INFORMATION: Synthetic Primer
401 <400> SEQUENCE: 15

```

VERIFICATION SUMMARY

DATE: 11/27/2001

PATENT APPLICATION: US/09/856,339A

TIME: 10:15:49

Input Set : A:\Sequence Listing.txt

Output Set: N:\CRF3\11212001\I856339A.raw

09856339-031501  
"05F80" 6EE95850

# SEQUENCE LISTING

<110> Isis Innovation Limited  
Wong, Luet  
Bell, Stephen  
Carmichael, Angus

<120> Process for Oxidising Terpenes

<130> P02196US0

<140> US 09/856,339

<141> 2001-05-18

<150> PCT/GB99/03873

<151> 1999-11-19

<150> GB 9825421.2

<151> 1998-11-19

<160> 23

<170> PatentIn version 3.0

<210> 1

<211> 414

<212> PRT

<213> Pseudomonas putida

<400> 1

Thr Thr Glu Thr Ile Gln Ser Asn Ala Asn Leu Ala Pro Leu Pro Pro  
1 5 10 15

His Val Pro Glu His Leu Val Phe Asp Phe Asp Met Tyr Asn Pro Ser  
20 25 30

Asn Leu Ser Ala Gly Val Gln Glu Ala Trp Ala Val Leu Gln Glu Ser  
35 40 45

Asn Val Pro Asp Leu Val Trp Thr Arg Cys Asn Gly Gly His Trp Ile  
50 55 60

Ala Thr Arg Gly Gln Leu Ile Arg Glu Ala Tyr Glu Asp Tyr Arg His  
65 70 75 80

Phe Ser Ser Glu Cys Pro Phe Ile Pro Arg Glu Ala Gly Glu Ala Tyr  
85 90 95

Asp Phe Ile Pro Thr Ser Met Asp Pro Pro Glu Gln Arg Gln Phe Arg  
100 105 110

Ala Leu Ala Asn Gln Val Val Gly Met Pro Val Val Asp Lys Leu Glu  
115 120 125

Asn Arg Ile Gln Glu Leu Ala Cys Ser Leu Ile Glu Ser Leu Arg Pro  
130 135 140

Gln Gly Gln Cys Asn Phe Thr Glu Asp Tyr Ala Glu Pro Phe Pro Ile  
145 150 155 160

Arg	Ile	Phe	Met	Leu	Leu	Ala	Gly	Leu	Pro	Glu	Glu	Asp	Ile	Pro	His
				165					170					175	
Leu	Lys	Tyr	Leu	Thr	Asp	Gln	Met	Thr	Arg	Pro	Asp	Gly	Ser	Met	Thr
			180					185					190		
Phe	Ala	Glu	Ala	Lys	Glu	Ala	Leu	Tyr	Asp	Tyr	Leu	Ile	Pro	Ile	Ile
		195					200					205			
Glu	Gln	Arg	Arg	Gln	Lys	Pro	Gly	Thr	Asp	Ala	Ile	Ser	Ile	Val	Ala
	210					215					220				
Asn	Gly	Gln	Val	Asn	Gly	Arg	Pro	Ile	Thr	Ser	Asp	Glu	Ala	Lys	Arg
225					230					235					240
Met	Cys	Gly	Leu	Leu	Leu	Val	Gly	Gly	Leu	Asp	Thr	Val	Val	Asn	Phe
				245					250					255	
Leu	Ser	Phe	Ser	Met	Glu	Phe	Leu	Ala	Lys	Ser	Pro	Glu	His	Arg	Gln
			260					265					270		
Glu	Leu	Ile	Glu	Arg	Pro	Glu	Arg	Ile	Pro	Ala	Ala	Cys	Glu	Glu	Leu
		275					280					285			
Leu	Arg	Arg	Phe	Ser	Leu	Val	Ala	Asp	Gly	Arg	Ile	Leu	Thr	Ser	Asp
	290					295					300				
Tyr	Glu	Phe	His	Gly	Val	Gln	Leu	Lys	Lys	Gly	Asp	Gln	Ile	Leu	Leu
305				310						315					320
Pro	Gln	Met	Leu	Ser	Gly	Leu	Asp	Glu	Arg	Glu	Asn	Ala	Cys	Pro	Met
				325					330					335	
His	Val	Asp	Phe	Ser	Arg	Gln	Lys	Val	Ser	His	Thr	Thr	Phe	Gly	His
			340					345					350		
Gly	Ser	His	Leu	Cys	Leu	Gly	Gln	His	Leu	Ala	Arg	Arg	Glu	Ile	Ile
		355					360					365			
Val	Thr	Leu	Lys	Glu	Trp	Leu	Thr	Arg	Ile	Pro	Asp	Phe	Ser	Ile	Ala
	370					375					380				
Pro	Gly	Ala	Gln	Ile	Gln	His	Lys	Ser	Gly	Ile	Val	Ser	Gly	Val	Gln
385					390					395					400
Ala	Leu	Pro	Leu	Val	Trp	Asp	Pro	Ala	Thr	Thr	Lys	Ala	Val		
				405					410						

```
<210> 2
<211> 1242
<212> DNA
<213> Pseudomonas putida
```

```

ttttccagcg agtgcgcgtt catccctcgt gaagccggcg aagcctacga cttcattccc 300
acctcgatgg atccgcccga gcagcgccag tttcgtgcgc tggccaacca agtgggttggc 360
atgccggtgg tggataagct ggagaaccgg atccaggagc tggcctgctc gctgatcgag 420
agcctgcgcc cgcaaggaca gtgcaacttc accgaggact acgccgaacc cttcccgata 480
cgcatcttca tgctgctcgc aggtctaccg gaagaagata tcccgcaactt gaaataccta 540
acggatcaga tgaccgcgtc ggatggcagc atgaccttcg cagaggccaa ggaggcgctc 600
tacgactatc tgataccgat catcgagcaa cgcaggcaga agccgggaac cgacgctatc 660
agcatcggtt ccaacggcca ggtcaatggg cgaccgatca ccagtgcga agccaagagg 720
atgtgtggcc tgttactggt cggcggcctg gatacgggtg tcaatttcct cagcttcagc 780
atggagttcc tggccaaaag cccggagcat cgccaggagc tgatcgagcg tcccagagcg 840
attccagccg cttgcgagga actactccgg cgcttctcgc tggttgcga tggccgcata 900
ctcacctccg attacgagtt tcatggcgtg caactgaaga aaggtgacca gatcctgcta 960
ccgcagatgc tgtctggcct ggatgagcgc gaaaacgcct gcccgatgca cgtcgacttc 1020
agtcgcaaaa aggtttcaca caccaccttt ggccacggca gccatctgtg ccttggccag 1080
cacctggccc gccgggaaat catcgtcacc ctcaaggaat ggctgaccag gattcctgac 1140
ttctccattg ccccggtgc ccagattcag cacaagagcg gcatcgtcag cggcgtgcag 1200
gcactccctc tggctctggga tccggcgact accaaagcgg ta 1242

```

```

<210> 3
<211> 1049
<212> PRT
<213> Bacillus megaterium

<400> 3

```

```

Met Thr Ile Lys Glu Met Pro Gln Pro Lys Thr Phe Gly Glu Leu Lys
1          5          10          15

Asn Leu Pro Leu Leu Asn Thr Asp Lys Pro Val Gln Ala Leu Met Lys
          20          25          30

Ile Ala Asp Glu Leu Gly Glu Ile Phe Lys Phe Glu Ala Pro Gly Arg
          35          40          45

Val Thr Arg Tyr Leu Ser Ser Gln Arg Leu Ile Lys Glu Ala Cys Asp
          50          55          60

Glu Ser Arg Phe Asp Lys Asn Leu Ser Gln Ala Leu Lys Phe Val Arg
65          70          75          80

Asp Phe Ala Gly Asp Gly Leu Phe Thr Ser Trp Thr His Glu Lys Asn
          85          90          95

```

105 FEB 66 0937

Trp	Lys	Lys	Ala	His	Asn	Ile	Leu	Leu	Pro	Ser	Phe	Ser	Gln	Gln	Ala	
			100					105					110			
Met	Lys	Gly	Tyr	His	Ala	Met	Met	Val	Asp	Ile	Ala	Val	Gln	Leu	Val	
		115					120					125				
Gln	Lys	Trp	Glu	Arg	Leu	Asn	Ala	Asp	Glu	His	Ile	Glu	Val	Pro	Glu	
	130					135					140					
Asp	Met	Thr	Arg	Leu	Thr	Leu	Asp	Thr	Ile	Gly	Leu	Cys	Gly	Phe	Asn	
145					150					155					160	
Tyr	Arg	Phe	Asn	Ser	Phe	Tyr	Arg	Asp	Gln	Pro	His	Pro	Phe	Ile	Thr	
			165						170					175		
Ser	Met	Val	Arg	Ala	Leu	Asp	Glu	Ala	Met	Asn	Lys	Leu	Gln	Arg	Ala	
		180						185					190			
Asn	Pro	Asp	Asp	Pro	Ala	Tyr	Asp	Glu	Asn	Lys	Arg	Gln	Phe	Gln	Glu	
		195					200					205				
Asp	Ile	Lys	Val	Met	Asn	Asp	Leu	Val	Asp	Lys	Ile	Ile	Ala	Asp	Arg	
	210					215					220					
Lys	Ala	Ser	Gly	Glu	Gln	Ser	Asp	Asp	Leu	Leu	Thr	His	Met	Leu	Asn	
225					230					235					240	
Gly	Lys	Asp	Pro	Glu	Thr	Gly	Glu	Pro	Leu	Asp	Asp	Glu	Asn	Ile	Arg	
				245					250					255		
Tyr	Gln	Ile	Ile	Thr	Phe	Leu	Ile	Ala	Gly	His	Glu	Thr	Thr	Ser	Gly	
		260						265					270			
Leu	Leu	Ser	Phe	Ala	Leu	Tyr	Phe	Leu	Val	Lys	Asn	Pro	His	Val	Leu	
		275					280					285				
Gln	Lys	Ala	Ala	Glu	Glu	Ala	Ala	Arg	Val	Leu	Val	Asp	Pro	Ala	Pro	
	290					295					300					
Ser	Tyr	Lys	Gln	Val	Lys	Gln	Leu	Lys	Tyr	Val	Gly	Met	Val	Leu	Asn	
305					310					315					320	
Glu	Ala	Leu	Arg	Leu	Trp	Pro	Thr	Ala	Pro	Ala	Phe	Ser	Leu	Tyr	Ala	
			325						330					335		
Lys	Glu	Asp	Thr	Val	Leu	Gly	Gly	Glu	Tyr	Pro	Leu	Glu	Lys	Gly	Asp	
		340						345					350			
Glu	Leu	Met	Val	Leu	Ile	Pro	Gln	Leu	His	Arg	Asp	Lys	Thr	Ile	Trp	
		355					360					365				
Gly	Asp	Asp	Val	Glu	Glu	Phe	Arg	Pro	Glu	Arg	Phe	Glu	Asn	Pro	Ser	
	370					375					380					
Ala	Ile	Pro	Gln	His	Ala	Phe	Lys	Pro	Phe	Gly	Asn	Gly	Gln	Arg	Ala	
385					390					395					400	
Cys	Ile	Gly	Gln	Gln	Phe	Ala	Leu	His	Glu	Ala	Thr	Leu	Val	Leu	Gly	
			405						410					415		
Met	Met	Leu	Lys	His	Phe	Asp	Phe	Glu	Asp	His	Thr	Asn	Tyr	Glu	Leu	



420										425					430				
Asp	Ile	Lys	Glu	Thr	Leu	Thr	Leu	Lys	Pro	Glu	Gly	Phe	Val	Val	Lys				
		435					440					445							
Ala	Lys	Ser	Lys	Lys	Ile	Pro	Leu	Gly	Gly	Ile	Pro	Ser	Pro	Ser	Thr				
	450					455					460								
Glu	Gln	Ser	Ala	Lys	Lys	Ala	Arg	Lys	Lys	Ala	Glu	Asn	Ala	His	Asn				
465					470					475					480				
Thr	Pro	Leu	Leu	Val	Leu	Tyr	Gly	Ser	Asn	Met	Gly	Thr	Ala	Glu	Gly				
				485					490					495					
Thr	Ala	Arg	Asp	Leu	Ala	Asp	Ile	Ala	Met	Ser	Lys	Gly	Phe	Ala	Pro				
			500					505					510						
Gln	Val	Ala	Thr	Leu	Asp	Ser	His	Ala	Gly	Asn	Leu	Pro	Arg	Glu	Gly				
		515					520					525							
Ala	Val	Leu	Ile	Val	Thr	Ala	Ser	Tyr	Asn	Gly	His	Pro	Pro	Asp	Asn				
	530					535					540								
Ala	Lys	Gln	Phe	Val	Asp	Trp	Leu	Asp	Gln	Ala	Ser	Ala	Asp	Glu	Val				
545					550					555					560				
Lys	Gly	Val	Arg	Tyr	Ser	Val	Phe	Gly	Cys	Gly	Asp	Lys	Asn	Trp	Ala				
				565					570					575					
Thr	Thr	Tyr	Gln	Lys	Val	Pro	Ala	Phe	Ile	Asp	Glu	Thr	Leu	Ala	Ala				
			580					585					590						
Lys	Gly	Ala	Glu	Asn	Ile	Ala	Asp	Arg	Gly	Glu	Ala	Asp	Ala	Ser	Asp				
		595					600					605							
Asp	Phe	Glu	Gly	Thr	Tyr	Glu	Glu	Trp	Arg	Glu	His	Met	Trp	Ser	Asp				
	610					615					620								
Val	Ala	Ala	Tyr	Phe	Asn	Leu	Asp	Ile	Glu	Asn	Ser	Glu	Asp	Asn	Lys				
625					630					635					640				
Ser	Thr	Leu	Ser	Leu	Gln	Phe	Val	Asp	Ser	Ala	Ala	Asp	Met	Pro	Leu				
				645					650					655					
Ala	Lys	Met	His	Gly	Ala	Phe	Ser	Thr	Asn	Val	Val	Ala	Ser	Lys	Glu				
			660					665					670						
Leu	Gln	Gln	Pro	Gly	Ser	Ala	Arg	Ser	Thr	Arg	His	Leu	Glu	Ile	Glu				
		675					680					685							
Leu	Pro	Lys	Glu	Ala	Ser	Tyr	Gln	Glu	Gly	Asp	His	Leu	Gly	Val	Ile				
	690					695					700								
Pro	Arg	Asn	Tyr	Glu	Gly	Ile	Val	Asn	Arg	Val	Thr	Ala	Arg	Phe	Gly				
705					710					715					720				
Leu	Asp	Ala	Ser	Gln	Gln	Ile	Arg	Leu	Glu	Ala	Glu	Glu	Glu	Lys	Leu				
				725					730					735					
Ala	His	Leu	Pro	Leu	Ala	Lys	Thr	Val	Ser	Val	Glu	Glu	Leu	Leu	Gln				
			740					745					750						

Tyr Val Glu Leu Gln Asp Pro Val Thr Arg Thr Gln Leu Arg Ala Met  
755 760 765

Ala Ala Lys Thr Val Cys Pro Pro His Lys Val Glu Leu Glu Ala Leu  
770 775 780

Leu Glu Lys Gln Ala Tyr Lys Glu Gln Val Leu Ala Lys Arg Leu Thr  
785 790 795 800

Met Leu Glu Leu Leu Glu Lys Tyr Pro Ala Cys Glu Met Lys Phe Ser  
805 810 815

Glu Phe Ile Ala Leu Leu Pro Ser Ile Arg Pro Arg Tyr Tyr Ser Ile  
820 825 830

Ser Ser Ser Pro Arg Val Asp Glu Lys Gln Ala Ser Ile Thr Val Ser  
835 840 845

Val Val Ser Gly Glu Ala Trp Ser Gly Tyr Gly Glu Tyr Lys Gly Ile  
850 855 860

Ala Ser Asn Tyr Leu Ala Glu Leu Gln Glu Gly Asp Thr Ile Thr Cys  
865 870 875 880

Phe Ile Ser Thr Pro Gln Ser Glu Phe Thr Leu Pro Lys Asp Pro Glu  
885 890 895

Thr Pro Leu Ile Met Val Gly Pro Gly Thr Gly Val Ala Pro Phe Arg  
900 905 910

Gly Phe Val Gln Ala Arg Lys Gln Leu Lys Glu Gln Gly Gln Ser Leu  
915 920 925

Gly Glu Ala His Leu Tyr Phe Gly Cys Arg Ser Pro His Glu Asp Tyr  
930 935 940

Leu Tyr Gln Glu Glu Leu Glu Asn Ala Gln Ser Glu Gly Ile Ile Thr  
945 950 955 960

Leu His Thr Ala Phe Ser Arg Met Pro Asn Gln Pro Lys Thr Tyr Val  
965 970 975

Gln His Val Met Glu Gln Asp Gly Lys Lys Leu Ile Glu Leu Leu Asp  
980 985 990

Gln Gly Ala His Phe Tyr Ile Cys Gly Asp Gly Ser Gln Met Ala Pro  
995 1000 1005

Ala Val Glu Ala Thr Leu Met Lys Ser Tyr Ala Asp Val His Gln  
1010 1015 1020

Val Ser Glu Ala Asp Ala Arg Leu Trp Leu Gln Gln Leu Glu Glu  
1025 1030 1035

Lys Gly Arg Tyr Ala Lys Asp Val Trp Ala Gly  
1040 1045

<210> 4  
<211> 3150  
<212> DNA

<213> Bacillus megaterium

<400> 4

atgacaatta aagaaatgcc tcagccaaaa acgtttggag agcttaaaaa ttaccgtta	60
ttaaacacag ataaaccggt tcaagctttg atgaaaattg cggatgaatt aggagaaatc	120
tttaaattcg aggcgcttg tctgtaatca cgctttgata aaaacttaag tcaagcgctt	180
aaatttgtac gtgtaacgag ctacttatca agtcagcgct taattaaaga agcatgcat	240
gattttgcag gagacgggtt atttacaagc tggacgcatg aaaaaaattg gaaaaagcg	300
cataatatct tacttccaag cttcagtcag caggcaatga aaggctatca tgcgatgatg	360
gtcgatatcg ccgtgcagct tgttcaaaaag tgggagcgct taaatgcaga tgagcatatt	420
gaagtaccgg aagacatgac acgtttaacg cttgatacaa ttggtctttg cggctttaac	480
tatcgcttta acagctttta ccgagatcag cctcatccat ttattacaag tatggtccgt	540
gcactggatg aagcaatgaa caagctgcag cgagcaaatc cagacgacc agcttatgat	600
gaaaacaagc gccagtttca agaagatatc aaggatgatga acgacctagt agataaaatt	660
attgcagatc gcaaagcaag cggatgaaca agcgatgatt tattaacgca tatgctaaac	720
ggaaaagatc cagaaacggg tgagccgctt gatgacgaga acattcgcta tcaaattatt	780
acattcttaa ttgcgggaca cgaaacaaca agtgggtctt tatcatttgc gctgtatttc	840
ttagtgaaaa atccacatgt attacaaaaa gcagcagaag aagcagcacg agttctagta	900
gatcctgctc caagctacaa acaagtcaaa cagcttaaat atgtcggcat ggtcttaaac	960
gaagcgctgc gcttatggcc aactgctcct gcgttttccc tatatgcaaa agaagatacg	1020
gtgcttggag gagaatatcc tttagaaaaa ggcgacgaac taatggttct gattcctcag	1080
cttcaccgtg ataaaacaat ttggggagac gatgtggaag agttccgtcc agagcgtttt	1140
gaaaatccaa gtgcgattcc gcagcatgag tttaaaccgt ttggaaacgg tcagcgctgcg	1200
tgtatcgggc agcagttcgc tcttcatgaa gcaacgctgg tacttggtat gatgctaaaa	1260
cactttgact ttgaagatca taaaaactac gagctggata ttaaagaaac tttaacgtta	1320
aaacctgaag gctttgtggt aaaagcaaaa tcgaaaaaaaa ttccgcttgg cggatttcct	1380
tcacctagca ctgaacagtc tgccaaaaaa gcacgcaaaa aggagaaaa cgctcataat	1440
acgcgcgtgc ttgtgctata cggttcaaat atgggaacag ctgaaggaac ggcgcgtgat	1500
ttagcagata ttgcaatgag caaaggattt gcaccgcagg tcgcaacgct tgattcacac	1560
gccggaaatc ttccgcgcga aggagctgta ttaattgtaa cggcgtctta taacggtcat	1620
ccgcctgata acgcaaagca atttgtcgac tggttagacc aagcgtctgc tgatgaagta	1680
aaaggcgttc gctactccgt atttggatgc ggcgataaaa actgggctac tacgtatcaa	1740

aaagtgcctg cttttatcga tgaaacgctt gccgctaaag gggcagaaaa catcgctgac 1800  
cgcggtgaag cagatgcaag cgacgacttt gaaggcacat atgaagaatg gcgtgaacat 1860  
atgtggagtg acgtagcagc ctactttaac ctcgacattg aaaacagtga agataataaa 1920  
tctactcttt cacttcaatt tgtcgacagc gccgcgata tgccgcttgc gaaaatgcac 1980  
ggtgcgtttt caacgaacgt cgtagcaagc aaagaacttc aacagccagg cagtgcacga 2040  
agcacgcgac atcttgaaat tgaacttcca aaagaagctt cttatcaaga aggagatcat 2100  
ttaggtgtta ttcttcgcaa ctatgaagga atagtaaacc gtgtaacagc aaggttcggc 2160  
ctagatgcat cacagcaa at cgtcttgaa gcagaagaag aaaaattagc tcatttgcca 2220  
ctcgctaaaa cagtatccgt agaagagctt ctgcaatacg tggagcttca agatcctgtt 2280  
acgcgcacgc agcttcgcg aatggctgct aaaacggtct gcccgccgca taaagtagag 2340  
cttgaagcct tgcttgaaaa gcaagcctac aaagaacaag tgctggcaaa acgtttaaca 2400  
atgcttgaac tgcttgaaaa ataccggcg tgtgaaatga aattcagcga atttatcgcc 2460  
cttctgcaa gcatacgccc gcgctattac tcgatttctt catcacctcg tgtcgatgaa 2520  
aaacaagcaa gcatcacggt cagcgttgtc tcaggagaag cgtggagcgg atatggagaa 2580  
tataaaggaa ttgctgcgaa ctatcttgcc gagctgcaag aaggagatac gattacgtgc 2640  
tttatttcca caccgcagtc agaatttacg ctgcaaaaag accctgaaac gccgcttato 2700  
atggtcggac cggaacagg cgtcgcgccg ttagaggct ttgtgcaggc gcgcaaacag 2760  
ctaaaagaac aaggacagtc acttgagaa gcacatttat acttcggctg ccgttcacct 2820  
catgaagact atctgtatca agaagagctt gaaaacgccc aaagcgaagg catcattacg 2880  
cttcataccg ctttttctcg catgccaaat cagccgaaaa catacgttca gcacgtaatg 2940  
gaacaagacg gcaagaaatt gattgaactt cttgatcaag gagcgcaact ctatatattgc 3000  
ggagacggaa gccaaatggc acctgccgtt gaagcaacgc ttatgaaaag ctatgctgac 3060  
gttcaccaag tgagtgaagc agacgctcgc ttatggctgc agcagctaga agaaaaaggc 3120  
cgatacgcaa aagacgtgtg ggctgggtaa 3150

<210> 5  
<211> 51  
<212> DNA  
<213> Artificial Sequence

<400> 5  
gagattaaga attcataaac acatgggagt gcgtgccata tgaacgcaa c 51

<210> 6  
<211> 15  
<212> PRT

<213> Artificial Sequence

<400> 6

Thr His Arg Ala Ser Pro Gly Leu Tyr Gly Leu Tyr Ser Glu Arg  
1 5 10 15

<210> 7

<211> 36

<212> DNA

<213> Artificial Sequence

<400> 7

gaactgagta gtgccactga cggaggatcc tcatcg 36

<210> 8

<211> 36

<212> DNA

<213> Artificial Sequence

<400> 8

cgatgaggat cctccgtcag tggcactact cagttc 36

<210> 9

<211> 6

<212> PRT

<213> Artificial Sequence

<400> 9

Gly Leu Tyr Ser Glu Arg  
1 5

<210> 10

<211> 36

<212> DNA

<213> Artificial Sequence

<400> 10

tcatcgggat cctcatcgat gtctaaagta gtgtat 36

<210> 11

<211> 52

<212> DNA

<213> Artificial Sequence

<400> 11

cccgataggc aatggtaatc atcgggagtc tagagcatcg aagctttcat cg 52

<210> 12

<211> 52

<212> DNA

<213> Artificial Sequence

<400> 12

cgatgaaagc ttcgatgctc tagactcccg atgattacca ttgcctatcg gg 52

<210> 13  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<400> 13  
 tcatcgacgc gtcgcgaact gctg 24

<210> 14  
 <211> 69  
 <212> DNA  
 <213> Artificial Sequence

<400> 14  
 cccgataggc aatggtaagt aggtgaatat ctaatcccca tctatgcgcg agtggagtct 60  
 agagttcga 69

<210> 15  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<400> 15  
 tcatcgaagc ttggctgttt t 21

<210> 16  
 <211> 47  
 <212> DNA  
 <213> Artificial Sequence

<400> 16  
 acaatttcac acaggatcta gaccatatgt catcgaagct ttcatcg 47

<210> 17  
 <211> 47  
 <212> DNA  
 <213> Artificial Sequence

<400> 17  
 cgatgaaagc ttcgatgaca tatggtctag atcctgtgtg aaattgt 47

<210> 18  
 <211> 21  
 <212> PRT  
 <213> Artificial sequence

<400> 18  
 Thr His Arg Ala Ser Pro Gly Leu Tyr Thr His Arg Ser Glu Arg Ser  
 1 5 10 15  
 Glu Arg Thr His Arg  
 20

<210> 19  
<211> 21  
<212> PRT  
<213> Artificial sequence

<400> 19

Thr His Arg Ala Ser Pro Gly Leu Tyr Ala Leu Ala Ser Glu Arg Ser  
1 5 10 15

Glu Arg Ser Glu Arg  
20

<210> 20  
<211> 17  
<212> PRT  
<213> Artificial sequence

<400> 20

Thr Asp Gly Thr Arg Pro Gly Pro Gly Pro Gly Pro Gly Pro Ser Ser  
1 5 10 15

Thr

<210> 21  
<211> 21  
<212> PRT  
<213> Artificial sequence

<400> 21

Thr Asp Gly Thr Arg Pro Gly Pro Gly Pro Gly Pro Gly Pro Gly Pro  
1 5 10 15

Gly Pro Ser Ser Thr  
20

<210> 22  
<211> 4  
<212> PRT  
<213> Artificial sequence

<400> 22

Pro Leu Glu Leu  
1

<210> 23  
<211> 21  
<212> PRT  
<213> Artificial sequence

<400> 23

Thr His Arg Ala Ser Pro Gly Leu Tyr Gly Leu Tyr Ser Glu Arg Ser  
1 5 10 15

Glu Arg Ser Glu Arg  
20

-54-

CLAIMS

1. Process for oxidising a substrate which is an acyclic or cyclic terpene, or a cycloalkene; or a substituted derivative thereof, which process comprises oxidising said compound with a mutant haem-containing enzyme, the mutant comprising the substitution of an amino acid in the active site by an amino acid with a less polar side-chain.

2. Process according to claim 1 in which the enzyme is a mutant of P450<sub>cam</sub> or P450<sub>BM-3</sub>, or a mutant of a naturally occurring homologue of either of these enzymes.

3. Process according to claim 2 in which the enzyme is one in which amino acid 47 and/or 51 of P450<sub>BM-3</sub>, or amino acid 96 of P450<sub>cam</sub>, or the equivalent amino acid(s) in a said homologue, have been changed to an amino acid with a less polar side-chain.

4. Process according to any one of claims 1, 2 or 3 in which there are one or more other amino acid substitutions in the active site.

5. Process according to any one of the preceding claims in which the enzyme is (i) P450<sub>cam</sub> and comprises one or more of the following mutations: F87W, F87I, F87L, T185L, T185F, V247A, V247L or F87A-I395F; or (ii) P450<sub>BM-3</sub> and comprises the mutation R47L-Y51F.

6. An enzyme as defined in claim 4 or 5 excluding mutants of P450<sub>cam</sub> which only have the mutations F87A-Y96G-F193A, F87A-Y96G-F193A-C334A, or T101M-T185F-V247M.

7. A polynucleotide which comprises a sequence which encodes an enzyme as defined in claim 6.

8. A cell which expresses:

(i) an enzyme as defined in any one of claims 2 to 6 which in its naturally occurring form has an electron transfer reductase domain, or

(ii) (a) an enzyme as defined in any one of claims 1 to

5;



-55-

(b) an electron transfer reductase; and

(c) an electron transfer redoxin.

9. A cell which expresses:

(a) (i) P450<sub>cam</sub>, or a fragment thereof; or

(ii) a naturally occurring homologue of P450<sub>cam</sub>  
or a fragment thereof; or

(iii) a mutant P450<sub>cam</sub>, or a mutant  
homologue of thereof  
as defined in any one of claims 2 to 5; or

(iv) a P450<sub>cam</sub> which has at least 70% amino acid  
homology with (i), (ii) or (iii) and  
optionally has the mutations defined in  
any one of claims 3 to 5; and

(b) an electron transfer reductase; and

(c) an electron transfer redoxin:

excluding an *E. Coli* DH5 $\alpha$  cell in which the only  
mutants of P450<sub>cam</sub> which are expressed are amongst the  
following:

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTSST-putidaredoxin reductase-TDGASSS-  
putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPSST-putidaredoxin  
reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-P450<sub>cam</sub>-TDGTRPGPGPGPGPGPSST-putidaredoxin  
reductase-TDGASSS-putidaredoxin-COOH,

H<sub>2</sub>N-putidaredoxin reductase-TDGASSS-putidaredoxin-  
PLEL-P450<sub>cam</sub>-COOH.

10. A cell according to claim 8 or 9 in which (a),  
(b) and (c) or (b) and (c) are expressed together in the  
same fusion protein.

11. A cell according to any one of claims 8 to 10  
in which:

(b) is putidaredoxin reductase or a fragment  
thereof; and/or

(c) is putidaredoxin or a fragment thereof.

-56-

12. A cell which expresses:

- (a) (i) P450<sub>BM-3</sub>, or a fragment thereof; or  
(ii) a naturally occurring homologue of P450<sub>BM-3</sub>  
or a fragment thereof; or  
(iii) a mutant P450<sub>BM-3</sub>, or a mutant  
homologue of thereof  
as defined in any one of claims 2 to 5.

13. A process according to any one of claims 1 to 5  
in which the compound is oxidised in a cell according to  
any one of claims 8 to 12

14. A process for making a library of mutants of  
P450<sub>cam</sub> or P450<sub>BM-3</sub>, or mutants of a homologue of either of  
these enzymes comprising contacting a cell according to  
any one of claims 9, and 10 and 11 when dependent on  
claim 9, or according to claim 12 or an *E.Coli* DH5 $\alpha$  cell  
as defined in claim 9;

with a mutagen and/or when then the cell is a  
mutator cell culturing the cell in conditions in which  
mutants are produced.

15. Process for selecting a mutant of P450<sub>cam</sub> or  
P450<sub>BM-3</sub>, or a homologue thereof, for its ability to  
oxidise a particular substrate, which process comprises  
screening a group of said mutants for their oxidation  
effect on the particular substrate.

16. Process according to claim 15 in which the  
mutant is additionally selected for its ability to  
oxidise the particular compound to a particular oxidation  
product.


17. A process according to claim 15 or 16 in which  
the screening is carried out on the library made in a  
process according to claim 14.

18. A process for producing a library of oxidation  
products comprising providing a substrate as defined in  
claim 1 to a library made in a process according to claim  
14 and allowing oxidation of the substrate.

-57-

19. An oxidation product obtained by a process according to any one of claims 1 to 5 and 13 wherein optionally the enzyme is one which has been selected in a process according to claim 15, 16 or 17 for use in a method of treatment of the human or animal body by therapy.

20. A pharmaceutical composition comprising an oxidation product obtained by a process according to any one of claims 1 to 5 and 13 wherein optionally the enzyme is one which has been selected in a process according to claim 15, 16 or 17 and a pharmaceutically acceptable carrier or diluent.






Fig.3.




Fig.4.

